arXiv:alg-geom/9602008v1 9 Feb 1996

Spin Verlinde spaces and Prym theta functions

W.M. Oxbury

Department of Mathematical Sciences
University of Durham

1 Introduction

The Verlinde formula is a remarkable—and potentially very useful—new tool in the geometry of algebraic curves which is borrowed from conformal field theory. In the first instance it is a trigonometric expression which assigns a natural number $N_l(G,g)$ to data consisting of a semisimple algebraic group G, a nonnegative integer g and an auxiliary integer $l \in \mathbf{Z}$. In physics $N_l(G,g)$ is interpreted as the dimension of the space of conformal blocks at level l in the Wess-Zumino-Witten model of conformal field theory on a compact Riemann surface of genus g. In algebraic geometry this 'space of conformal blocks' is identified with a vector space of the form $H^0(\mathcal{M}_C(G), \mathcal{L}^l)$, where $\mathcal{M}_C(G)$ is the moduli scheme (or stack) of semistable principal G-bundles over C, and $\mathcal{L}$ is an ample line bundle on this moduli space generating the Picard group.

The feature of the Verlinde formula which motivates this paper is its 'numerology'. Namely, when one computes the numbers $N_l(G,g)$ for the classical simple groups one finds that they obey interesting identities which lead one to certain conjectures about the geometry of $\mathcal{M}_C(G)$. (See [OW].) In this article we are concerned with identities linking the complex spin groups $G = \mathrm{Spin}_m$ with the configuration of principally polarised Prym varieties associated to the curve C via its unramified double covers. This connection was first observed in [O2] for the odd spin groups Spin_{2n+1}; here we shall give a systematic account of both the odd and even cases.

The moduli space $\mathcal{M}_C(SO_m)$ has, for $m \geq 3$, two connected components labelled by the second Stiefel-Whitney class w_2 of the bundles. Each of these components has a $J_2(C)$-Galois cover which is a moduli space for Clifford bundles with fixed spinor norm line bundle $\xi \in \mathrm{Pic}(C)$, and whose isomorphism class depends only on $\deg \xi$ mod 2. When $\xi = \mathcal{O}_C$ the Galois cover is precisely $\mathcal{M}_C(\mathrm{Spin}_m)$ since by definition Spin_m is the kernel of the spinor norm; the 'sister' space which arises when $\deg \xi$ is odd, we denote by $\mathcal{M}_C^-(\mathrm{Spin}_m)$. In each case the fibre $J_2(C)$ over $\mathcal{M}_C(SO_m)$ parametrises liftings of a given SO_m-bundle to a Clifford bundle with given spinor norm ξ with $\deg \xi \equiv w_2$ mod 2.

Note that for low values of m we recover well-known moduli spaces of vector bundles: for example $\mathcal{M}_C(\mathrm{Spin}_3) = \mathcal{SU}_C(2,0)$ and $\mathcal{M}_C^-(\mathrm{Spin}_3) = \mathcal{SU}_C(2,1)$ (where $\mathcal{SU}_C(n,d)$ is the moduli space of rank n vector bundles with fixed determinant of degree d); while $\mathcal{M}_C(\mathrm{Spin}_6) = \mathcal{SU}_C(4,0)$ and $\mathcal{M}_C^-(\mathrm{Spin}_6) = \mathcal{SU}_C(4,2)$.

We shall consider the theta line bundle $\Theta(\mathbf{C}^m)$ on these varieties, coming from the standard orthogonal representation $\mathbf{C}^m$ of the Clifford group, and use the Verlinde formula to count its sections.

These preliminary ideas—the spin moduli spaces, theta line bundles on them and the Verlinde calculations—occupy the first five sections of the paper. The central observations on which the paper is based is contained in section 6. This is that the dimension of $H^0(\mathcal{M}_C(\mathrm{Spin}_m), \Theta(\mathbf{C}^m))$ coincides with that of the direct sum of the spaces of *even* level m theta functions on all the Prym varieties (including the Jacobian itself); while the dimension of $H^0(\mathcal{M}_C^-(\mathrm{Spin}_m), \Theta(\mathbf{C}^m))$ is equal to that of the direct sum of the spaces of *odd* level m theta functions on the Prym varieties. The precise statement (see theorem 6.1 and table (22)) depends on the parity of m: when m is even we have to take theta functions not just on the Prym varieties P_η, $\eta \in J_2(C)\backslash\{\mathcal{O}\}$, but on the two component abelian subvarieties

$$P_\eta \cup P_\eta^- = \mathrm{Nm}^{-1} K_C \subset J^{2g-2}(\widetilde{C}_\eta),$$

where $\widetilde{C}_\eta \to C$ is the double cover corresponding to each η.

There are two remarks to make about this feature of the even spin groups. The first is that it is natural in the sense that, whereas level m theta functions are well-defined on P_η for all m—there is a canonically defined theta divisor Ξ_η—on P_η^- they are well-defined only if m is even. The second is that it corresponds to a direct sum decomposition of $H^0(\mathcal{M}_C(\mathrm{Spin}_m), \Theta(\mathbf{C}^m))$ into two pieces when m is even—apparently the eigenspaces under the involution of the moduli space $\mathcal{M}_C(\mathrm{Spin}_m)$ corresponding to reflection of the Dynkin diagram. (See section 7.1.)

In the remaining sections of the paper we make sense of the observations of section 6 by constructing homomorphisms:

$$H^0(\mathcal{M}^{\pm}(\mathrm{Spin}_{2n+1}), \Theta(\mathbf{C}^{2n+1}))^{\vee} \to \sum_{\eta \in J_2} H^0_{\pm}(P_\eta, (2n+1)\Xi_\eta)$$

$$\begin{aligned} H^0(\mathcal{M}^{\pm}(\mathrm{Spin}_{2n}), \Theta(\mathbf{C}^{2n}))^{\vee} \quad \to \quad & \sum_{\eta \in J_2} H^0_{\pm}(P_\eta, 2n\Xi_\eta) \\ & \oplus \sum_{\eta \neq 0} H^0_{\pm}(P^-_\eta, 2n\Xi_\eta). \end{aligned}$$

Table (22) asserts that each of these maps is *between vector spaces of equal dimension*. It is necessary to emphasise that the validity of the right-hand column of the table is dependent on a natural-seeming conjecture 5.7 for a Verlinde formula on $\mathcal{M}^-(\mathrm{Spin}_m)$. When $m = 3$ this is the Verlinde formula for rank 2 vector bundles of odd degree due to Thaddeus [T]; further evidence for this 'twisted' formula is given in [OW].

It is therefore natural to expect that the above maps are isomorphisms; for $m \geq 5$ this is not known, though we hope to return to the question in a later paper. Cases of low m, on the other hand, where the spin moduli spaces can be identified with more familiar moduli spaces of vector bundles, are examined individually in section 6.

Finally, when m is odd it is known (see [LS]) that $\Theta(\mathbf{C}^m) = 2\mathcal{P}$ where the 'Pfaffian' line bundle $\mathcal{P}$ generates the Picard group. In section 6.1 we observe—though this remark is independent of the rest of the paper—that the space of sections of this line bundle has a basis labelled by the even theta characteristics of the curve, which directly generalises that constructed by Beauville in [B2] for the case $m = 3$.

Acknowledgements: In writing this paper the author has benefited greatly from conversations with B. van Geemen, C. Pauly, S. Ramanan and C. Sorger, to all of whom he expresses his gratitude.

2 Moduli spaces of principal bundles on a curve

In this section we shall give a brief account of the moduli spaces of semistable principal bundles over a curve, following [R1], [DN], [KNR].

We begin with a smooth projective complex curve C of genus $g \geq 2$, and a complex connected reductive algebraic group G; and we consider algebraic principal G-bundles $E \to C$. Topologically such bundles are classified by the fundamental group of G.

Just as for vector bundles, one has notions of stability, semistability and S-equivalence for algebraic G-bundles, and for stable bundles S-equivalence is the same as isomorphism. (We shall recall in a moment the definition of stability, but it will not be necessary here to define S-equivalence.) The basic result of Ramanathan [R2] is then the following.

2.1 Theorem. *Given C, G as above and an element $\gamma \in \pi_1(G)$, there exists a normal irreducible projective variety $\mathcal{M}(G,\gamma)$ which is a coarse moduli space for families of semistable G-bundles of type γ on C, modulo S-equivalence.*

Moreover, one has

$$\dim \mathcal{M}(G,\gamma) = (g-1)\dim G + \dim Z(G),$$

and $\mathcal{M}(G,\gamma)$ is unirational when G is a simple group [KNR].

The basic construction with principal bundles is the following. If E is a G-bundle, and $\rho : G \to \mathrm{Aut}(X)$ any left G-space, then we can form a bundle $E(X) = E \times_\rho X$ with fibre X. In case $X = G/P$ is a homogeneous coset space, a section $\sigma : C \to E(G/P)$ is called a reduction of the strucure group of the bundle to the subgroup P. When $P \subset G$ is a maximal parabolic, $E(G/P) \to C$ can be thought of as a 'generalised Grassmannian bundle'. Then by definition, E is *semistable* if and only if

$$\deg \sigma^* T^{\mathrm{vert}} E(G/P) \geq 0 \qquad \textit{for all maximal parabolics } P \subset G,$$

where T^{vert} denotes the vertical tangent bundle.

On the other hand, if $\pi : G' \to G$ is a group epimorphism then we can view $X = G$ as a left G'-space via π, and so form a G-bundle $E = F(G)$ from any G'-bundle F. F is said to be a *lift* of E. In particular, if G' is a central extension of G then there is a bijection between maximal parabolics $P \subset G$ and maximal parabolics $P' = \pi^{-1}P \subset G'$, and moreover $F(G'/P') \cong E(G/P)$ if F is any lift of E. Consequently:

2.2 Lemma. *If E is a G-bundle and F a lift of E to a central extension of G then E is stable (resp. semistable) if and only if F is.*

Finally, of course, we can take for the G-space X a finite-dimensional representation $\rho : G \to GL(V)$, to obtain a vector bundle $E(V)$. In the case when $G = GL_n$ and $V = \mathbf{C}^n$ is the standard representation, the notions of stability, semistability and S-equivalence are the same for the principal bundle E as for the vector bundle $E(V)$. Thus we shall write $\mathcal{U}(n,d) = \mathcal{M}(GL_n, d)$, for $d \in \pi_1(GL_n) \cong \mathbf{Z}$; this is the moduli space of semistable vector bundles of rank n and degree d.

Consider now the determinant morphism

$$\det : \mathcal{U}(n,d) \to J^d(C).$$

This is a fibration and we shall, as is usual, denote the isomorphism class of the fibre by $\mathcal{SU}(n,d) = \mathcal{SU}_C(n,d)$; or $\mathcal{SU}_C(n)$ when $d = 0$.

One knows from [DN] that, via det, $\mathcal{U}(n,d)$ has Picard group

$$\text{Pic } \mathcal{U}(n,d) \cong \text{Pic } J^d(C) \oplus \mathbf{Z}\{\Theta_{n,d}\}$$

where $\Theta_{n,d}$ is an ample line bundle on the fibres constructed as follows. It will be convenient, to begin with, to assume that n divides d, i.e. that we are dealing with vector bundles of integral slope.

Consider first an arbitrary family $F \to C \times S$ of semistable vector bundles on C with rank n, degree d and slope $\mu = d/n \in \mathbf{Z}$, as above; and we construct a line bundle $\Theta(F) \to S$, functorial with respect to base change $S' \to S$, in the following way. Let $\pi : C \times S \to S$ be the projection. Then (at least Zariski locally) there is a homomorphism of locally free sheaves on S, $\phi : K^0 \to K^1$, having the direct images of F under π as kernel and cokernel:

$$0 \to R^0_\pi F \to K^0 \xrightarrow{\phi} K^1 \to R^1_\pi F \to 0. \tag{1}$$

Moreover, the determinant line bundle

$$\text{Det}(F) = (\textstyle\bigwedge^{\text{top}} K^0)^\vee \otimes (\textstyle\bigwedge^{\text{top}} K^1)$$

is well-defined and functorial with respect to base change. If $\mu = g-1$ we write $\Theta(F) = \mathrm{Det}(F)$; and this has a canonical section $\det\phi$, so that in this case $\Theta(F)$ is represented by a canonical Cartier divisor on S. Otherwise $\Theta(F)$ is defined to be a suitable twist of $\mathrm{Det}(F)$ such that

$$\Theta(F \otimes \pi^* L) = \Theta(F)$$

for any line bundle $L \to S$; i.e. Θ respects equivalence of families.

Now in the case $\mu = g-1$ it is shown in [DN] that the functor Θ is in fact represented in moduli space by a global Cartier divisor

$$\Theta_{n,n(g-1)} = \mathrm{Closure}\{\text{stable } V | H^0(V) = H^1(V) \neq 0\} \subset \mathcal{U}(n, n(g-1)).$$

In other words $\Theta(F) = f^*\Theta_{n,n(g-1)}$ where $f : S \to \mathcal{U}(n, n(g-1))$ is given by the coarse moduli property.

For the general case ($\mu \in \mathbf{Z}$ still) one chooses a line bundle $L \in \mathrm{Pic}(C)$ with degree chosen so that we get a morphism

$$\mathcal{U}(n,d) \xrightarrow{\otimes L} \mathcal{U}(n, n(g-1)).$$

Now set

$$\Theta_L = (\otimes L)^* \Theta_{n,n(g-1)}.$$

The dependence of Θ_L on L is then given by (2) below, which is a consequence of:

2.3 Lemma. *View $J^0(C)$ as a subgroup of* $\mathrm{Pic}(J^d(C))$ *by $L \mapsto \Phi^{-1} \otimes T_L^*\Phi$, where Φ is any line bundle representing the principal polarisation on $J^d(C)$. Then for any family $F \to C \times S$ as above, and any $L \in J^0(C)$, we have*

$$\Theta(F \otimes pr_C^* L) = \det{}^*(L) \otimes \Theta(F)$$

where $\det : S \to \mathcal{U}(n,d) \to J^d(C)$.

It follows easily from this that when L, L' have the same degree, Θ_L and $\Theta_{L'}$ are related by:

$$\Theta_{L'} = \det{}^*(L'L^{-1}) \otimes \Theta_L \in \mathrm{Pic}\,\mathcal{U}(n,d). \tag{2}$$

We now set $\Theta_{n,d} = \Theta_L$: if $d \neq n(g-1)$ this depends on L, *but by (2) its restriction to the fibres of* $\det : \mathcal{U}(n,d) \to J^d(C)$ *is independent of L.*

In order to consider bundles of general degree, i.e. non-integral slope, it is necessary to twist by bundles L of higher rank:

$$\mathcal{U}(n,d) \xrightarrow{\otimes L} \mathcal{U}(rn, rn(g-1)),$$

where L has rank r. It is easy to check that the necessary and sufficient condition for arranging slope $g-1$ on the right is that:

$$r \in \frac{n}{\gcd(n,d)}\mathbf{Z} \tag{3}$$

The line bundle Θ_L may now be defined in the same way as above. In this more general situation (2) becomes:

$$\Theta_{L'} = \det^*(\det L' \otimes \det L^\vee) \otimes \Theta_L \in \operatorname{Pic} \mathcal{U}(n,d), \tag{4}$$

Consequently the restriction of Θ_L to the fibres of $\det : \mathcal{U}(n,d) \to J^d(C)$ is again independent of the choice of L with given rank r; and we set $\Theta_{n,d} = \Theta_L$ for any L with $r = n/\gcd(n,d)$. This is the required generator of the Picard group.

Note that if in this construction L, L' are two vector bundles of different ranks $r < r'$ (both satisfying (3), and the degrees of L and L' chosen suitably) then

$$\Theta_{L'} = \Theta_L^{\otimes r'/r}. \tag{5}$$

Finally, suppose that we are given a family $E \to C \times S$ of semistable G-bundles, and a representation $\rho : G \to SL(V)$, where $\dim V = n$. We shall suppose that ρ satisfies the condition:

$$Z(G)_0 \subset \ker \rho \tag{6}$$

We can form the family of vector bundles $E(V) \to C \times S$; and by [Rth2] proposition 2.17 the condition (6) guarantees that these vector bundles are semistable.

We thus obtain a theta line bundle $\Theta(E(V)) \to S$, and since $E(V)$ has trivial determinant on the fibres of $\pi : C \times S \to S$ we deduce from lemma 2.3 the following corollary, which will be needed later:

2.4 Corollary. *For $E \to C \times S$ and $\rho : G \to SL(V)$ as above, and for any $L \in J^0(C)$ one has*

$$\Theta(E(V) \otimes pr_C^* L) = \Theta(E(V)).$$

Globally ρ satisfying (6) induces a morphism

$$\rho_* : \mathcal{M}(G,\gamma) \to \mathcal{M}(SL_n) \hookrightarrow \mathcal{U}(n,0)$$

and the functor $E \mapsto \Theta(E(V))$ is represented by the (well-defined) line bundle

$$\Theta(V) := (\rho_*)^*\Theta_{n,0} \in \mathrm{Pic}\ \mathcal{M}(G,\gamma).$$

Note that if we let $j : \mathcal{M}(SL_n) \hookrightarrow \mathcal{U}(n,0)$ denote the inclusion which identifies $\mathcal{M}(SL_n)$ with the moduli space of vector bundles of rank n and trivial determinant, via the standard representation $\mathbf{C}^n$, then by construction

$$\Theta(\mathbf{C}^n) = j^*\Theta_{n,0}.$$

3 Clifford bundles

Let us consider again the fibration

$$\det : \mathcal{U}_C(n,d) \to J^d(C);$$

induced, that is, by the determinant homomorphism $GL_n \to \mathbf{C}^*$. The fibres of these maps are, up to isomorphism, the n moduli varieties $\mathcal{SU}_C(n,d)$, for $d \in \mathbf{Z}/n$. In this section we shall describe an alternative generalisation of this situation for $n = 2$, obtained by replacing GL_2 not by GL_n, but by the special Clifford group of a nondegenerate quadratic form. First we need to recall some basic Clifford theory.

3.1 The special Clifford group

Let Q be a nondegenerate quadratic form on a complex vector space V of finite dimension m; let $A = A(Q)$ be its Clifford algebra and A^+ the even Clifford algebra. Recall that these can be expressed as matrix algebras as follows.

If $m = 2n$ is even then for any n-dimensional isotropic subspace $U \subset V$ one has

$$A \cong \mathrm{End} \textstyle\bigwedge U; \qquad A^+ \cong \mathrm{End}(\textstyle\bigwedge^{\mathrm{even}} U) \oplus \mathrm{End}(\textstyle\bigwedge^{\mathrm{odd}} U). \tag{7}$$

If, on the other hand, $m = 2n+1$ is odd then for any direct sum decomposition $V = U \oplus U' \oplus \mathbf{C}$ where U, U' are n-dimensional isotropic subspaces one has

$$A \cong \operatorname{End} \textstyle\bigwedge U \oplus \operatorname{End} \textstyle\bigwedge U'; \qquad A^+ \cong \operatorname{End} \textstyle\bigwedge U. \tag{8}$$

The 'principal involution' of A is $\alpha : x \mapsto -x$ for $x \in V$, i.e. is ± 1 on $A^\pm$ respectively. The 'principal anti-involution' β is the identity on V and reverses the direction of multiplication: $\beta(x_1 \dots x_r) = x_r \dots x_1$. Then the *Clifford group* is

$$C(Q) = \{s \in A^* | \alpha(s) V s^{-1} \subset V\},$$

where $A^* \subset A$ denotes the group of units; and the *special Clifford group* is

$$SC(Q) = C(Q) \cap A^+.$$

For $s \in C(Q)$ the transformation $\pi_s : x \mapsto \alpha(s) x s^{-1}$ of V is orthogonal—this is because $C(Q)$ is generated by $x \in V \cap C(Q)$, for which π_x is just minus the reflection in the hyperplane $x^\perp$. Thus one has a group homomorphism $\pi : C(Q) \to O(Q)$, which has the following properties.

3.1 Proposition. *1.* $\ker \pi = \mathbf{C}^*$*;*

2. $\pi(C(Q)) = O(Q)$ *and* $\pi(SC(Q)) = SO(Q)$*.*

3.2 Corollary. $SC(Q)$ *is a connected reductive algebraic group.*

The *spinor norm* is the group homomorphism

$$\begin{array}{rccl} \operatorname{Nm} : & SC(Q) & \to & \mathbf{C}^* \\ & s & \mapsto & \beta(s)s. \end{array}$$

Equivalently $\operatorname{Nm}(x_1 \dots x_r) = Q(x_1) \cdots Q(x_r)$ for $x_1, \dots, x_r \in V$. Then by definition $\operatorname{Spin}(Q) = \ker \operatorname{Nm}$.

Note that multiplication by scalars induces a double cover

$$\{\pm(1,1)\} \to \mathbf{C}^* \times \operatorname{Spin}(Q) \to SC(Q). \tag{9}$$

From now on we shall write C_m, SC_m, Spin_m instead of $C(Q)$, $SC(Q)$, $\operatorname{Spin}(Q)$ when Q is the standard quadratic form on $\mathbf{C}^m$. Then (using (3.1)) one has the following commutative diagram of short exact sequences:

$$\begin{tikzcd}
1 \arrow[r] & \mathbf{C}^* \arrow[r] & SC_m \arrow[r, "\pi"] & SO_m \arrow[r] & 1 \\
(10) & & & & \\
1 \arrow[r] & \mathbf{Z}/2 \arrow[r] \arrow[uu] & \mathrm{Spin}_m \arrow[r] \arrow[uu] & SO_m \arrow[r] \arrow[uu, equal] & 1.
\end{tikzcd}$$

3.3 Proposition. *For $m \geq 3$:*

1. *SC_m has centre $Z(SC_m) = \begin{cases} \mathbf{C}^* & \text{if } m \text{ is odd} \\ \mathbf{C}^* \times \mathbf{Z}/2 & \text{if } m \text{ is even;} \end{cases}$*
2. *SC_m has fundamental group $\pi_1(SC_m) = \mathbf{Z}$; and this maps isomorphically to $\pi_1(\mathbf{C}^*) = \mathbf{Z}$ under the spinor norm.*

Proof. (i) If m is odd the centre of SC_m must be contained in—and hence equal to—the kernel $\mathbf{C}^*$ of the surjection onto SO_m, since the latter has trivial centre.

If, on the other hand, m is even, then by the same token $Z(SC_m)$ is contained in $\pi^{-1}(Z(SO_m))$ where π denotes the surjection to SO_m. In this case SO_m has centre $\{\pm 1\}$. As before everything in $\pi^{-1}(1) = \mathbf{C}^*$ is central; while if $\{e_1, \ldots, e_m\} \subset \mathbf{C}^m$ is any orthonormal basis then the product $e_1 \ldots e_m \in SC_m$ spans $\pi^{-1}(-1) \cong \mathbf{C}^*$. Since m is even this product anticommutes with each e_i, and therefore *commutes* with all elements of A^+. So $\pi^{-1}\{\pm 1\} \cong \mathbf{C}^* \times \mathbf{Z}/2$ is contained in and therefore equal to the centre.

(ii) From the exact homotopy sequence of the fibration in the upper sequence of (10), and the vanishing of $\pi_2(\mathbf{C}^*)$, we have a non-split extension

$$0 \to \mathbf{Z} \to \pi_1(SC_m) \to \mathbf{Z}/2 \to 0.$$

Since the fundamental group of a Lie group is abelian it follows that the only possibility is $\pi_1(SC_m) = \mathbf{Z}$. The last part now follows from the fact that Spin_m is simply-connected. □

3.2 The spin moduli spaces

From proposition 3.3 part 2 we see that there is for each $d \in \mathbf{Z} = \pi_1(SC_m)$ a morphism induced by the spinor norm, and which we shall denote in the same way:

$$\mathrm{Nm} : \mathcal{M}(SC_m, d) \to J^d(C).$$

Moreover, when $m = 3$ this is nothing but the determinant morphism for rank 2 vector bundles (see example 3.7 below). And just as for rank 2 vector bundles, one has:

3.4 Proposition. *For $d \in \mathbf{Z}$ and $L \in J^d(C)$, the isomorphism class of the subscheme* $\mathrm{Nm}^{-1}(L) \subset \mathcal{M}(SC_m, d)$ *depends only on d mod 2.*

Actually this is essentially trivial and is proved in the same way as for rank 2 vector bundles, once one observes that multiplication of SC_m by its centre (proposition 3.3) induces a natural generalisation of the tensor product operation of Clifford bundles by line bundles if m is odd, and if m is even by *pairs* (N, η) where N is a line bundle and $\eta \in H^1(C, \mathbf{Z}/2) = J_2(C)$. (See also [R1].) We shall write, respectively, $N \otimes E$ and $(N, \eta) \otimes E$ for this product. It follows from the definition of the spinor norm that

$$\mathrm{Nm}(N \otimes E) = N^2 \otimes \mathrm{Nm}(E), \quad \text{resp.} \quad \mathrm{Nm}((N, \eta) \otimes E) = N^2 \otimes \mathrm{Nm}(E).$$

So to prove 3.4: suppose $d = \deg L \equiv d' = \deg L'$ mod 2, and write $L' = N^{2k} \otimes L$ for some $N \in \mathrm{Pic}^1(C)$, $k \in \mathbf{Z}$. Then the map $\mathcal{M}(SC_m, d) \to \mathcal{M}(SC_m, d')$ given by $E \mapsto N \otimes E$ (resp. $(N, \mathcal{O}) \otimes E$) restricts to an isomorphism $\mathrm{Nm}^{-1}(L) \cong \mathrm{Nm}^{-1}(L')$. $\square$

We shall therefore introduce the notation:

$$\begin{array}{rcl} \mathcal{M}^+(\mathrm{Spin}_m) & = & \mathcal{M}(\mathrm{Spin}_m) = \mathrm{Nm}^{-1}(\mathcal{O}_C); \\ \mathcal{M}^-(\mathrm{Spin}_m) & = & \mathrm{Nm}^{-1}(\mathcal{O}_C(p)), \end{array}$$

where $p \in C$ is any point of the curve.

3.5 *Remark.* The group SO_m has fundamental group $\mathbf{Z}/2$; so the moduli space of semistable SO_m-bundles has two irreducible components $\mathcal{M}(SO_m, 0) = \mathcal{M}^+(SO_m)$ and $\mathcal{M}(SO_m, 1) = \mathcal{M}^-(SO_m)$ distinguished by the second Stiefel-Whitney class w_2. It is not hard to show (see [O2]) that $\deg \mathrm{Nm}(E) \equiv w_2(E)$ mod 2 and that $\mathcal{M}^{\pm}(\mathrm{Spin}_m)$ are naturally Galois covers of these components:

$$\mathcal{M}^{\pm}(\mathrm{Spin}_m) \xrightarrow{J_2(C)} \mathcal{M}^{\pm}(SO_m).$$

Note that these maps respect stability—this is a special case of lemma 2.2.

In later sections we shall be interested in the theta line bundle $\Theta(\mathbf{C}^m)$ associated to the orthogonal representation of SC_m. Note that by 3.1 and 3.3 the condition (6) is satisfied for $m \geq 3$, so that $\Theta(\mathbf{C}^m)$ is defined everywhere on $\mathcal{M}(SC_m)$ and hence on both $\mathcal{M}^{\pm}(\mathrm{Spin}_m)$.

3.6 Example. m = 2. Here $SC_2 \cong \mathbf{C}^* \times \mathbf{C}^*$, the spinor norm is $\mathrm{Nm} : (a,b) \mapsto ab$ and the orthogonal representation $SC_2 \to SO_2 \cong \mathbf{C}^*$ is $(a,b) \mapsto a/b$. It follows that $\mathcal{M}(SC_2) \cong \mathrm{Pic}(C) \times \mathrm{Pic}(C)$ and each of $\mathcal{M}^{\pm}(\mathrm{Spin}_2) \cong \mathrm{Pic}(C)$.

$\Theta(\mathbf{C}^2)$ is by definition obtained by pulling back $\Theta_{2,0}$ under

$$\rho_* : \mathcal{M}(SC_2) = \mathrm{Pic}(C) \times \mathrm{Pic}(C) \to \mathcal{M}(SL_2).$$

But this map sends a pair of line bundles (L, N) to the vector bundle $LN^{-1} \oplus L^{-1}N$, and this is semistable only if $\deg L = \deg N$. (Note that condition (6) fails for this case!) It follows that the morphism ρ_*, and hence $\Theta(\mathbf{C}^2)$, is defined only on $\mathrm{Pic}(C) \times_{\deg} \mathrm{Pic}(C)$; and $\Theta(\mathbf{C}^2)$ is thus defined only on the degree 0 component $J(C)$ of $\mathcal{M}(\mathrm{Spin}_2) = \mathrm{Pic}(C)$ and is not defined on $\mathcal{M}^-(\mathrm{Spin}_2)$.

On the other hand, $\mathcal{M}(\mathrm{Spin}_2) \to \mathcal{M}(SO_2)$ is the squaring map [2] on line bundles, and so one sees that on $J(C)$ the orthogonal theta bundle is:

$$\Theta(\mathbf{C}^2) = [2]^*(2\theta) = 8\theta,$$

where θ is the theta divisor on $J(C)$.

3.7 Example. m = 3. This is in many ways the most important case. It is well-known that SC_3 is, via (8), equal to the group of units $GL_2 \subset A^+$, and that the spinor norm is the determinant homomorphism so that $\mathrm{Spin}_3 = SL_2$. (The representation on SO_3 is then precisely the action of GL_2 on $S^2 \cong \mathbf{P}_1$ by Möbius transformations, via stereographic projection.) Thus $\mathcal{M}(\mathrm{Spin}_3) \cong \mathcal{SU}_C(2)$ and $\mathcal{M}^-(\mathrm{Spin}_3) \cong \mathcal{SU}_C(2,1)$.

Let $\mathcal{L}_d = \Theta_{2,d}$ be the (ample) generators of the Picard groups of these varieties, for $d = 0, 1$ respectively, as described in the previous section. From the discussion there we can view $\mathcal{L}_1$ ($= \Theta_{L'}$ for a suitable rank 2 vector bundle L') as defined on all of $\mathcal{M}(GL_2)$ while $\mathcal{L}_0$ ($= \Theta_L$ for a suitable line bundle L) is a line bundle defined only on the components of even degree; and by (5) $\mathcal{L}_1 = \mathcal{L}_0^2$ on these components.

The orthogonal theta bundle is then $\Theta(\mathbf{C}^3) = \mathcal{L}_1^2 = \mathcal{L}_0^4$—see (19) below.

3.8 Example. m = 4. Again it is very well-known that $\mathrm{Spin}_4 = SL_2 \times SL_2$. Via (7) the special Clifford group $SC_4 \subset GL_2 \times GL_2$ is the subgroup consisting of pairs of matrices (A, B) such that $\det A = \det B$; the spinor norm is then the common 2×2 determinant. Thus $\mathcal{M}(SC_4) = \mathcal{M}(GL_2) \times_{\det} \mathcal{M}(GL_2)$ while $\mathcal{M}^{\pm}(\mathrm{Spin}_4)$ are $\mathcal{SU}_C(2) \times \mathcal{SU}_C(2)$ and $\mathcal{SU}_C(2,1) \times \mathcal{SU}_C(2,1)$ respectively.

The orthogonal representation induces $\mathcal{M}(SC_4) \to \mathcal{M}(SL_4)$ mapping a pair of rank 2 vector bundles (E, F) to $E \otimes F^*$, and from this it follows that the orthogonal theta bundle is (with the notation of the previous example)

$$\Theta(\mathbf{C}^4) = pr_+^* \mathcal{L}_1 \otimes pr_-^* \mathcal{L}_1 = pr_+^* \mathcal{L}_0^2 \otimes pr_-^* \mathcal{L}_0^2,$$

where $pr_{\pm}$ denote the respective projections.

3.9 Example. m = 6. In this case $\mathrm{Spin}_6 \cong SL_4$, and one may show that the subgroup (using (7) once again) $SC_6 \subset GL_4 \times GL_4$ is the image of the homomorphism

$$\begin{array}{rcl} \mathbf{C}^* \times SL_4 & \to & GL_4 \times GL_4 \\ (\lambda, A) & \mapsto & (\lambda A, \lambda A^{adj,t}) \end{array}$$

where $A^{adj,t}$ is the matrix of signed cofactors and $A^{adj} = \det A \times A^{-1} = A^{-1}$. (Compare with (9).)

Note that projection to the first factor—the first half-spinor representation—induces a double cover $\{(1, \pm 1)\} \subset SC_6 \xrightarrow{pr_1} GL_4$. Moreover, the lift of the determinant function to this double cover has a square root—namely, the spinor norm $\mathrm{Nm} : (\lambda A, \lambda A^{adj,t}) \mapsto \lambda^2$, where $\mathrm{Nm}(a)^2 = \det pr_1(a)$ for $a \in SC_6$.

At the level of bundles this says we have a commutative diagram

$$\begin{tikzcd} \mathcal{M}(SC_6) \arrow[r, "pr_1"] \arrow[d, "\mathrm{Nm}"'] & \mathcal{M}(GL_4) \arrow[d, "\det"] \\ \mathrm{Pic}(C) \arrow[r, "{[2]}"] & \mathrm{Pic}(C) \end{tikzcd}$$

where each of the horizontal maps has fibre $J_2(C)$. In particular one sees that $\mathcal{M}(\mathrm{Spin}_6) \cong \mathcal{SU}_C(4)$ and $\mathcal{M}^-(\mathrm{Spin}_6) \cong \mathcal{SU}_C(4,2)$.

As in example 3.7, let $\mathcal{L}_d = \Theta_{2,d}$ be the ample generators of the Picard groups of these varieties, for $d = 0, 2$ respectively. Then $\mathcal{L}_2$ ($= \Theta_{L'}$ for a suitable rank 2 vector bundle L') is defined on all components of $\mathcal{M}(GL_2)$ of even degree, while $\mathcal{L}_0$ ($= \Theta_L$ for a suitable line bundle L) is defined only on the components of degree $\equiv 0 \bmod 4$; and by (5) $\mathcal{L}_2 = \mathcal{L}_0^2$ on these components. Using (19) from section 5 below, the orthogonal theta bundle is then $\Theta(\mathbf{C}^6) = \mathcal{L}_2 = \mathcal{L}_0^2$.

4 Orthogonal bundles and theta characteristics

In this section we shall gather together various properties of orthogonal bundles, some possibly well-known, which will be needed later on.

4.1 Isotropic line subbundles

To begin, consider any SO_m-bundle E and its associated orthogonal vector bundle $E(\mathbf{C}^m)$. Recall that stability of E is equivalent to the condition that $\deg F < 0$ for all *isotropic* vector subbundles $F \subset E(\mathbf{C}^m)$. This holds, in particular, if $E(\mathbf{C}^m)$ is stable as a vector bundle.

On the other hand, for *any* subbundle $F \subset E(\mathbf{C}^m)$, the direct sum with its orthogonal complement fits into an exact sequence

$$0 \to N \to F \oplus F^\perp \to M \to 0$$

where N, M are the subbundles generically generated by $F \cap F^\perp$ and $F + F^\perp$ respectively. In particular, when $N = 0$ this gives rise to an orthogonal splitting $E(\mathbf{C}^m) = F \oplus F^\perp$. Applying this idea inductively Ramanan shows ([R1] proposition 4.5):

4.1 Lemma. *E is a stable SO_m-bundle if and only if $E(\mathbf{C}^m)$ is an orthogonal direct sum $E(\mathbf{C}^m) = F_1 \oplus \cdots \oplus F_k$ where the F_i are pairwise non-isomorphic stable vector bundles.*

In particular $E(\mathbf{C}^m)$ is a stable vector bundle for generic $E \in \mathcal{M}(SO_m)$.

(The last assertion here follows from a simple dimension count.)

Now suppose that $F \subset E(\mathbf{C}^m)$ is a *line* subbundle. Then $N = 0$ precisely when F is non-isotropic, and thus $E(\mathbf{C}^m)$ splits in this case, and so fails to be stable as a vector bundle. This shows:

4.2 Corollary. *For generic* $E \in \mathcal{M}(SO_m)$ *every line subbundle* $F \subset E(\mathbf{C}^m)$ *is isotropic.*

In the next lemma we observe that the bound $\deg F < 0$ for isotropic subbundles of $E(\mathbf{C}^m)$—again restricting to line subbundles—can in fact be improved generically:

4.3 Lemma. *For generic stable* $E \in \mathcal{M}^{\pm}(SO_m)$ *every isotropic line subbundle* $F \subset E(\mathbf{C}^m)$ *satisfies* $\deg F \leq -g+1$.

Proof. We simply count dimensions of those bundles $E \in \mathcal{M}(SO_m)$ which can possess an isotropic line subbundle $F \subset E(\mathbf{C}^m)$ with $\deg F = d$. Let $F^{\perp} \subset E(\mathbf{C}^m)$ denote the orthogonal complement: this is a vector bundle of rank $m-1$ on which the quadratic form restricts with rank $m-2$, and the quotient $Q = F^{\perp}/F$ is thus an SO_{m-2}-bundle, fitting into an exact diagram:

$$\begin{tikzcd}
 & 0 \arrow[d] & 0 \arrow[d] & & \\
 & F \arrow[r, equal] \arrow[d] & F \arrow[d] & & \\
0 \arrow[r] & F^{\perp} \arrow[r] \arrow[d] & E(\mathbf{C}^m) \arrow[r] \arrow[d] & F^{-1} \arrow[r] \arrow[d, equal] & 0 \\
0 \arrow[r] & Q \arrow[r] \arrow[d] & (F^{\perp})^{\vee} \arrow[r] \arrow[d] & F^{-1} \arrow[r] & 0 \\
 & 0 & 0 & &
\end{tikzcd}$$

Noting now that E is determined up to isomorphism by $F^{\perp}$ together with its (degenerate) quadratic form (which determines F), we see that to construct such a bundle E it is enough to specify the left-hand vertical sequence. For this, F is determined by g parameters; Q by $(g-1)(m-2)(m-3)/2$ parameters; and the extension $F^{\perp}$ by

$$h^1(C, Q^{\vee} \otimes F) - 1 = (m-2)(g-1-d) - 1$$

parameters. Consequently, a *generic* $E \in \mathcal{M}(SO_m)$ (in either component of the moduli space) possesses an isotropic line subbundle F of degree d only if the total number of parameters is at least $\dim \mathcal{M}(SO_m) = (g-1)m(m-1)/2$:

$$\begin{aligned} g + (g-1)(m-2)(m-3)/2 & \\ +(m-2)(g-1-d) - 1 \quad &\geq \quad (g-1)m(m-1)/2, \end{aligned}$$

which simplifies to $d \leq -g+1$. The lemma follows at once from this. □

4.4 *Remark.* Note that in the case $m = 3$, $E(\mathbf{C}^3) = \mathrm{ad}\ V$, the bundle of tracefree endomorphisms of a stable rank 2 vector bundle V—i.e. that coming from E via the 2-dimensional spin representation. Then there is a one-to-one correspondence between line subbundles $L \subset V$ and isotropic line subbundles $F \subset \mathrm{ad}\ V$: this is because the quadratic form on ad V is the Killing form, which for tracefree 2×2 matrices is the determinant. Thus F is isotropic if and only if it consists of nilpotent endomorphisms, and L is then the kernel bundle; and conversely $F = \mathrm{Hom}(V/L, L) = L^2 \otimes \det V^\vee$.

If one views $L \subset V$ as determining a cross-section $l \subset \mathbf{P}(V)$ of the corresponding ruled surface, then $-\deg F = l \cdot l$ is its self-intersection. So lemma 4.3 says

$$l \cdot l \geq g - 1 \qquad \text{for any section } l \subset \mathbf{P}(V),$$

for a generic rank 2 vector bundle V. A well-known result of Nagata [N], on the other hand, says that *every* ruled surface has a section l with self-intersection $l \cdot l \leq g$. Since, for a given surface, self-intersection of a section is constant ($\equiv \deg V$) mod 2, this implies that the inequality of lemma 4.3 is sharp for $m = 3$.

4.2 Conservation of parity

We shall need to make use of the following well-known result of Atiyah [A] and Mumford [M]:

4.5 Atiyah-Mumford lemma. *Suppose that $F \to C$ is a vector bundle admitting a nondegenerate symmetric bilinear form*

$$F \otimes F \to K_C.$$

Then $h^0(C, F)$ is constant modulo 2 under deformation.

The case of this which we shall be interested in arises when $F = L \otimes E(\mathbf{C}^m)$ for L a theta characteristic, $L^2 = K$, and E an SO_m- (or SC_m-) bundle. Let $\vartheta(C) \subset J^{g-1}(C)$ denote the set of theta characteristics. The value of $h^0(C, F)$ mod 2 is in this case given by a calculation of Serre:

4.6 Proposition. *For any SC_m-bundle $E \to C$ and theta characteristic $K^{\frac{1}{2}} \in \vartheta(C)$ one has*

$$h^0(C, K^{\frac{1}{2}} \otimes E(\mathbf{C}^m)) \equiv m h^0(C, K^{\frac{1}{2}}) + \deg \mathrm{Nm}(E) \mod 2.$$

Proof. By [S], theorem 2, one has the congruence:

$$h^0(C, K^{\frac{1}{2}} \otimes E(\mathbf{C}^m)) \equiv (m+1) h^0(C, K^{\frac{1}{2}}) + h^0(C, K^{\frac{1}{2}} \otimes w_1(E)) + w_2(E) \mod 2,$$

where w_1 and w_2 are the Stiefel-Whitney classes. But $w_1(E)$ can be identified with $\det E(\mathbf{C}^m) \in J_2(C) \cong H^1(C, \mathbf{Z}/2)$, which in our case vanishes since E is a *special* Clifford bundle. On the other hand, $w_2(E) \equiv \deg \mathrm{Nm}(E)$ mod 2 by remark 3.5. So we get the statement in the proposition. $\square$

4.7 Corollary. *Suppose that $p : \widetilde{C} \to C$ is an unramified double cover, and let $\sigma : \widetilde{C} \leftrightarrow \widetilde{C}$ denote the sheet-interchange over C. Suppose that E is any SC_m-bundle on C, and that $L \in \mathrm{Pic}(\widetilde{C})$ satisfies $L \otimes \sigma(L) = K_{\widetilde{C}}$. Then:*

$$h^0(\widetilde{C}, L \otimes p^* E(\mathbf{C}^m)) \equiv m h^0(\widetilde{C}, L) \mod 2.$$

Proof. L has degree $g(\widetilde{C}) - 1$, so by the Atiyah-Mumford lemma it suffices to assume that L is the pull-back of a theta characteristic. Then the corollary follows at once from the proposition since $w_2(p^* E) = 0$. $\square$

4.3 Generically zero results

The main aim of this section is to establish that in each of the results 4.6 and 4.7 above, the vanishing (mod 2) of the right-hand side guarantees that $h^0(C, L \otimes E(\mathbf{C}^m)) = 0$ for generic stable orthogonal bundle E (theorems 4.10, 4.12 and corollary 4.13). To this end we shall apply Brill-Noether methods (see, for example, [ACGH]) to the loci:

$$\mathcal{W}_k = \{(L, E) | h^0(C, L \otimes E(\mathbf{C}^m)) \geq k\} \subset J^{g-1}(C) \times \mathcal{N}^{st}(m)$$

where $\mathcal{N}^{st}(m) \subset \mathcal{N}(m)$ is the open set of *stable* bundles. Analogously with classical Brill-Noether theory we claim that the Zariski tangent space to $\mathcal{W}_k$ at a point (L, E) is annihilated by the image of a *Petri map*

$$\begin{aligned} \mu : H^0(L \otimes E(\mathbf{C}^m)) \otimes H^0(KL^{-1} \otimes E(\mathbf{C}^m)) \quad &\rightarrow \quad H^0(K) \oplus H^0(K \otimes E(\mathbf{so}_m)) \\ &\cong \Omega^1_{J^{g-1} \times \mathcal{N}^{st}(m)}|_{(L,E)}. \end{aligned}$$

Here $\mathbf{so}_m$ is the Lie algebra of Spin_m, viewed as the semisimple component of the adjoint representation of SC_m. So $E(\mathbf{so}_m)$ is then the vector bundle with fibre $\mathbf{so}_m$ associated to any Clifford bundle E via this representation; and by standard deformation theory (and this is where we require our bundles to be stable) $H^0(C, K \otimes E(\mathbf{so}_m)) \cong \Omega^1_{\mathcal{N}^{st}(m)}|_E$.

The Petri map is defined by

$$\mu : s \otimes t \mapsto \langle s, t \rangle \oplus s \wedge t \tag{11}$$

where $\langle , \rangle$ is the symmetric bilinear form on the vector bundle $E(\mathbf{C}^m)$, and we identify $\mathbf{so}_m \cong \bigwedge^2 \mathbf{C}^m$.

4.8 Proposition. *For $(L, E) \in \mathcal{W}_k - \mathcal{W}_{k+1}$ the Zariski tangent space of $\mathcal{W}_k$ at (L, E) is $T_{(L,E)}\mathcal{W}_k = (\text{image } \mu)^\perp$.*

To prove this we first need:

4.9 Lemma. *Given $(L, E) \in \mathrm{Pic}(C) \times \mathcal{N}^{st}(m)$, a section $s \in H^0(C, L \otimes E(\mathbf{C}^m))$, and a tangent vector $\eta \oplus \xi \in H^1(C, \mathcal{O}) \oplus H^1(C, E(\mathbf{so}_m)) = T_{(L,E)}(J^{g-1} \times \mathcal{N}^{st}(m))$, s extends to the 1st-order deformation of $L \otimes E(\mathbf{C}^m)$ corresponding to $\eta \oplus \xi$ if and only if*

$$\xi s + \eta s = 0 \in H^1(C, L \otimes E(\mathbf{C}^m)).$$

Proof. Represent E by transition data $\{h_{\alpha\beta}\}$ with respect to an open cover $\{U_\alpha\}$ of C, where the $h_{\alpha\beta}$ are holomorphic SC_m-valued functions on $U_\alpha \cap U_\beta$ satisfying the cocycle condition; let $g_{\alpha\beta}$ be the image of $h_{\alpha\beta}$ in SO_m—these are then the transition functions for the vector bundle $E(\mathbf{C}^m)$. Likewise represent L by transition data $\{\phi_{\alpha\beta}\}$ where the $\phi_{\alpha\beta}$ are $\mathbf{C}^*$-valued functions. Finally, represent the sum $\eta \oplus \xi \in H^1(\mathcal{O}) \oplus H^1(\mathbf{so}_m)$ by a cocycle $\{\eta_{\alpha\beta} \oplus \xi_{\alpha\beta}\}$ (holomorphic Lie algebra-valued functions on the $U_\alpha \cap U_\beta$).

Then the 1st-order deformation corresponding to $\eta \oplus \xi$ is the SO_m-bundle on $C \times \operatorname{Spec} \mathbf{C}[\varepsilon]/(\varepsilon^2)$ with transition data

$$\begin{aligned} \tilde{\phi}_{\alpha\beta} &= (1 + \varepsilon\eta_{\alpha\beta})\phi_{\alpha\beta}, \\ \tilde{g}_{\alpha\beta} &= (1 + \varepsilon\xi_{\alpha\beta})g_{\alpha\beta}. \end{aligned}$$

A section $s \in H^0(C, L \otimes E(\mathbf{C}^m))$ is now given by a collection $\{s_\alpha\}$ of $\mathbf{C}^m$-valued functions satisfying

$$s_\alpha = \phi_{\alpha\beta} g_{\alpha\beta} s_\beta$$

on $U_\alpha \cap U_\beta$; and s extends to the 1st-order deformation ξ provided there exists a 0-cochain $\{s'_\alpha\}$ such that

$$\tilde{s}_\alpha = s_\alpha + \varepsilon s'_\alpha$$

satisfies the cocycle condition

$$\begin{aligned} \tilde{s}_\alpha &= \tilde{\phi}_{\alpha\beta}\tilde{g}_{\alpha\beta}\tilde{s}_\beta \\ &= \phi_{\alpha\beta}(1 + \varepsilon\eta_{\alpha\beta})(1 + \varepsilon\xi_{\alpha\beta})g_{\alpha\beta}(s_\beta + \varepsilon s'_\beta) \\ &= s_\alpha + \varepsilon(\eta_{\alpha\beta}s_\alpha + \xi_{\alpha\beta}s_\alpha + \phi_{\alpha\beta}g_{\alpha\beta}s'_\beta), \end{aligned}$$

and hence $s'_\alpha - \phi_{\alpha\beta}g_{\alpha\beta}s'_\beta = (\eta_{\alpha\beta} + \xi_{\alpha\beta})s_\alpha$. In other words $s' = \{s'_\alpha\}$ has to satisfy

$$\eta s + \xi s = \partial s' \in Z^1(L \otimes E(\mathbf{C}^m)),$$

where ∂ is the coboundary operator on 0-cocycles, and Z^1 is the group of Cech 1-cocycles. So there exists a solution if and only if $\xi s + \eta s = 0$ in cohomology. □

Proof of proposition 4.8. We write down the Serre duality pairing of $\eta \oplus \xi \in H^1(C, \mathcal{O}) \oplus H^1(C, E(\mathbf{so}_m))$ with $\mu(s \otimes t)$, for $s \in H^0(C, L \otimes E(\mathbf{C}^m))$ and $t \in H^0(C, KL^{-1} \otimes E(\mathbf{C}^m))$:

$$\begin{aligned} (\eta \oplus \xi, \mu(s \otimes t))_{\text{Serre}} &= \int_C (\eta\langle s, t\rangle + \operatorname{trace}(\xi\ s \wedge t)) \\ &= \int_C \langle \xi s + \eta s, t\rangle, \end{aligned} \tag{12}$$

where we have identified $\bigwedge^2 \mathbf{C}^m$ with $\mathbf{so}_m$, the space of skew-symmetric $m \times m$ matrices, and used the fact that under this identification one has $\operatorname{trace}(\xi\ s \wedge t) = \langle \xi s, t\rangle$, as one verifies by an easy calculation.

Now for $(L,E) \in \mathcal{W}_k - \mathcal{W}_{k+1}$, the Zariski tangent space $T_{(L,E)}\mathcal{W}_k$ is the linear span of directions $\eta \oplus \xi$ in which *all* sections s extend. By the lemma this condition on $\eta \oplus \xi$ is that $\xi s + \eta s = 0$ in cohomology for all $s \in H^0(C, L \otimes E(\mathbf{C}^m))$. By (12) this linear span is precisely (image $\mu)^\perp$. $\square$

We come now to the first main result of this section:

4.10 Theorem. *For generic* $(L,E) \in J^{g-1}(C) \times \mathcal{M}^{\pm}(SO_m)$ *one has* $H^0(C, L \otimes E(\mathbf{C}^m)) = 0$.

Proof. We choose a component of $J^{g-1} \times \mathcal{N}^{st}(m)$, and suppose that $h^0(C, L \otimes E(\mathbf{C}^m)) \geq k > 0$ everywhere in this component, and generically equal to k, for some natural number k. We then choose a generic point (L,E) in this component; according to proposition 4.8 the Petri map μ is identically zero here. This means that for arbitrary sections $s \in H^0(C, L \otimes E(\mathbf{C}^m))$ and $t \in H^0(C, KL^{-1} \otimes E(\mathbf{C}^m))$ (and note that both spaces have the same dimension k, by Riemann-Roch) we have, on the one hand, $s \wedge t = 0$. This implies that s, t generically generate the same line subbundle $F \subset E(\mathbf{C}^m)$; moreover, since s, t are arbitrary we see that F is independent of their choice, and depends only on L and E.

On the other hand, again by (11), $\langle s, t \rangle = 0$; this implies that F is an *isotropic* line subbundle. (Of course, since E is generic this is also forced by corollary 4.2.) Lemma 4.3 and genericity of E therefore implies that $\deg F \leq -g+1$. But by definition of F the spaces of sections $H^0(L \otimes F)$ and $H^0(KL^{-1} \otimes F)$ are both nonzero, which forces $L \otimes F = KL^{-1} \otimes F = \mathcal{O}$ and hence $L^2 = K$. But this contradicts the genericity of L. $\square$

We now consider the intersection of the subscheme $\mathcal{W}_k$ with the fibre $\{K^{\frac{1}{2}}\} \times \mathcal{N}^{st}(m)$, where $K^{\frac{1}{2}} \in \vartheta(C)$ is a theta characteristic. (Indeed, theorem 4.10 above also follows—for all cases except $\mathcal{M}^-(SO_{2n})$—from theorem 4.12 below.)

Fixing $K^{\frac{1}{2}}$, one sees that by Serre duality the Petri map now factorises into two maps:

$$\begin{array}{rcl} \mu_J : S^2 H^0(C, K^{\frac{1}{2}} \otimes E(\mathbf{C}^m)) & \rightarrow & H^0(C, K), \\ \mu_{\mathcal{N}} : \bigwedge^2 H^0(C, K^{\frac{1}{2}} \otimes E(\mathbf{C}^m)) & \rightarrow & H^0(C, K \otimes \bigwedge^2 E(\mathbf{C}^m)) \\ & & \cong H^0(C, K \otimes E(\mathbf{so}_m)), \end{array}$$

where $\mu_J : s \otimes t \mapsto \langle s, t \rangle$, and $\mu_{\mathcal{N}}$ is the natural multiplication map.

As an immediate consequence of proposition 4.8 we obtain:

4.11 Corollary. *If* $E \in \mathcal{N}^{st}(m)$ *and* $K^{\frac{1}{2}} \in \vartheta(C)$ *satisfy* $h^0(C, K^{\frac{1}{2}} \otimes E(\mathbf{C}^m)) = k$ *then the subscheme* $\mathcal{U}_k = \mathcal{W}_k|_{\{K^{1/2}\}\times\mathcal{N}^{st}(m)} \subset \mathcal{N}^{st}(m)$ *has Zariski tangent space* $T_E\mathcal{U}_k = (\text{image } \mu_{\mathcal{N}})^{\perp}$.

From this follows our second main result:

4.12 Theorem. *For* $m \geq 3$ *and for any theta characteristic* $K^{\frac{1}{2}} \in \vartheta(C)$ *we have* $h^0(C, K^{\frac{1}{2}} \otimes E(\mathbf{C}^m)) = 0$ *or 1 for generic* $E \in \mathcal{M}^{\pm}(SO_m)$, *where the parity is determined by lemma 4.6.*

Proof. We fix our component $\mathcal{M}^{\pm}(SO_m)$, and suppose that $h^0(C, K^{\frac{1}{2}} \otimes E(\mathbf{C}^m)) \geq k$ everywhere in this component, generically equal to k. We then choose a generic point E in this component; by proposition 4.11 the Petri map $\mu_{\mathcal{N}}$ vanishes here. As in the proof of theorem 4.10 this means $s \wedge t = 0$ for all sections $s, t \in H^0(C, K^{\frac{1}{2}} \otimes E(\mathbf{C}^m))$, so all sections generate the same line subbundle $F \subset E(\mathbf{C}^m)$; in particular

$$H^0(C, K^{\frac{1}{2}} \otimes F) = H^0(C, K^{\frac{1}{2}} \otimes E(\mathbf{C}^m)).$$

Since E is generic it follows by corollary 4.2 that F is isotropic, and hence by lemma 4.3 that $\deg F \leq -g+1$. But then $\deg K^{\frac{1}{2}} \otimes F \leq 0$ and hence $k = h^0(C, K^{\frac{1}{2}} \otimes F) \leq 1$. □

4.13 Corollary. *Suppose that* $p : \widetilde{C} \to C$ *is an unramified double cover and that* E, L *is a generic pair as in corollary 4.7. Then—except possibly in the case of* m *even,* $w_2(E) \equiv 1$*—we have* $h^0(\widetilde{C}, L \otimes p^*E(\mathbf{C}^m)) = 0$ *or 1, where the parity is determined by corollary 4.7.*

Proof. As in the proof of 4.7 we take $L = p^*K^{\frac{1}{2}}$ to be the pull-back of a theta characteristic. (See also the discussion of section 7.2.) Then

$$h^0(\widetilde{C}, L \otimes p^*E(\mathbf{C}^m)) = h^0(C, K^{\frac{1}{2}} \otimes E(\mathbf{C}^m)) + h^0(C, K^{\frac{1}{2}} \otimes \eta \otimes E(\mathbf{C}^m)),$$

where $\eta \in J_2(C)$ is the 2-torsion point associated to the covering. If E is generic then by theorem 4.12 it is possible to choose $K^{\frac{1}{2}}$—in all cases except when m is even and $w_2(E)$ is odd—so that the right-hand side is ≤ 1. □

4.14 *Remark.* In the case of m even, $w_2(E)$ odd, $h^0(\widetilde{C}, L \otimes p^*E(\mathbf{C}^m)) \geq 2$ for any $L = p^*K^{\frac{1}{2}}$; so the above argument fails. Nonetheless, we expect that the result 4.13 is still true in this case, but its proof requires a refinement of the Brill-Noether analysis of this section.

5 The Verlinde formula

In this section we shall write down, for the unitary and spin groups, the Verlinde formula which calculates the dimension of the vector spaces $H^0(\mathcal{M}(G), \Theta(V))$. For the derivations of these formulae we refer the reader to [B3], [OW]. In fact, what one writes down is a natural number $N_l(G)$ depending on the group, on the genus g, and on an integer l called the 'level'. Then to any representation V of G, one associates a level $l = d_V$ such that $\dim H^0(\mathcal{M}(G), \Theta(V)) = N_{d_V}(G)$.

5.1 Preliminaries

For $k \in \mathbf{N}$ and $r \in \mathbf{Q}$ we let

$$f_k(r) = 4\sin^2(r\pi/k) = (1-\zeta_k^r)(1-\zeta_k^{-r})$$

where $\zeta_k = e^{2\pi i/k}$. This satisfies certain obvious identities (we shall usually drop the subscript k for convenience):

5.1 Lemma.

(i) $f(r) = f(-r)$;

(ii) $f(r) = f(k-r)$;

(iii) $f(k/2) = 4$;

(iv) $f(2r) = f(r)f(k/2-r)$;

(v) $\prod_{r=1}^{k-1} f(r) = k^2$.

In addition, we shall need the following:

5.2 Lemma.

(i) If $k = 2n+1$ *is odd then*

$$\begin{array}{rcl} \prod_{r=1}^{n} f(r) & = & 2n+1, \\ \prod_{r=1}^{2n} f(r)^{[\frac{r}{2}]} & = & (2n+1)^n. \end{array}$$

(ii) If $k = 2n$ *is even then*

$$\begin{array}{rcl} \prod_{r=1}^{n-1} f(r) & = & n, \\ \prod_{r=1}^{2n-1} f(r)^{[\frac{r}{2}]} & = & 2^{n-1}n^n \\ \prod_{r=1}^{2n-1} f(r)^{[\frac{r+1}{2}]} & = & 2^{n+1}n^n. \end{array}$$

Proof. (i) The first identity follows at once from parts (ii) and (v) of the previous lemma. For the second, use 5.1 (ii) to write

$$\prod_{r=1}^{2n} f(r)^{[\frac{r}{2}]} = \prod_{r=1}^{n} f(r)^{[\frac{r}{2}]+[\frac{2n+1-r}{2}]} = \prod_{r=1}^{n} f(r)^n = (2n+1)^n.$$

(ii) Again the first identity is an easy consequence of the previous lemma. We shall just give the proof of the third identity, that of the second being being almost the same.

The left-hand product can be rewritten, using 5.1 (ii), as:

$$P = f(n)^{[\frac{n+1}{2}]} \times \prod_{r=1}^{n-1} f(r)^{[\frac{r+1}{2}]+[\frac{2n-r+1}{2}]}.$$

We observe that

$$\Big[\frac{r+1}{2}\Big] + \Big[\frac{2n-r+1}{2}\Big] = \begin{cases} n & \text{if } r \text{ is even,} \\ n+1 & \text{if } r \text{ is odd.} \end{cases}$$

So suppose first that n is even. Using 5.1 (iii) and $\prod_{r=1}^{n-1} f(r) = n$ we see that

$$\begin{array}{rcl} P & = & 2^n f(1)^{n+1} f(2)^n \cdots f(n-2)^n f(n-1)^{n+1} \\ & = & (2n)^n f(1) f(3) \cdots f(n-1). \end{array}$$

We claim that $f(1)f(3)\cdots f(n-1) = 2$—from which the third identity in 5.2 (ii) follows. To see this, observe first that (using 5.1 (iv))

$$\begin{array}{rcl} f(2)f(4)\cdots f(n-2) & = & f(n/2)^{-1} \prod_{r=1}^{n-1} f(r) \\ & = & n/2; \end{array}$$

and second that

$$n = f(1)\cdots f(n-1) \times f(2)\cdots f(n-2).$$

The reasoning for n odd is similar. □

In what follows we shall consider sets $U = \{u_1, \ldots, u_n\}$ of rational numbers; and for such a set we define:

$$\begin{array}{rcl} \Psi_k(U) & = & \displaystyle\prod_{1\le i<j\le n} f(u_i - u_j) \\ \Pi_k(U) & = & \displaystyle\prod_{1\le i<j\le n} f(u_i - u_j) f(u_i + u_j) \\ \Phi_k(U) & = & \displaystyle\prod_{1\le i<j\le n} f(u_i - u_j) f(u_i + u_j) \times \prod_{i=1}^{n} f(u_i). \end{array}$$

5.3 Example. $\mathbf{SL_n}$. The Verlinde number in this case is:

$$N_l(SL_n) = \sum_U \Big(\frac{n(l+n)^{n-1}}{\Psi_{l+n}(U)}\Big)^{g-1}$$

where the sum is taken over $U = \{0 = u_0 < u_1 < \cdots < u_{n-1} < l+n\}$.

Since we are concerned with the spin groups we shall record at this point the result of computing this expression at levels 1 and 2 for $\mathrm{Spin}_6 = SL_4$:

$$\begin{array}{lrcl} & \bullet \ N_1(SL_4) & = & 2^{2g}; \\ (13) & & & \\ & \bullet \ N_2(SL_4) & = & 2^{3g-1}3^{g-1} + 2^{3g-1} + 2^g 3^{g-1}. \end{array}$$

5.4 Example. $\mathbf{Spin_{2n}}$, $\mathbf{n \geq 4}$. The Verlinde number at level l is here:

$$N_l(\mathrm{Spin}_{2n}) = \sum_{U \in P_l(2n)} \Big(\frac{4k^n}{\Pi_k(U)}\Big)^{g-1}, \qquad \text{where } k = l + 2n - 2;$$

where $P_l = P_l(2n)$ denotes the collection of sets $U = \{u_1 < \cdots < u_n\}$ satisfying the following conditions. (Note that this collection is finite provided $n \geq 3$.)

1. $u_i \in \frac{1}{2}\mathbf{Z}$;
2. $u_{i+1} - u_i \in \mathbf{Z}$ for $i = 1, \ldots n-1$;
3. $u_1 + u_2 > 0$; and
4. $u_{n-1} + u_n < k = l + 2n - 2$.

It will be convenient to write $P_l = P_l^+ \cup P_l^-$ where P_l^+ (resp. P_l^-) consists of U with all $u_i \in \mathbf{Z}$ (resp. $u_i \in \frac{1}{2}\mathbf{Z}\backslash\mathbf{Z}$). Correspondingly the Verlinde number splits up as $N_l = N_l^+ + N_l^-$ where

$$N_l^{\pm}(\mathrm{Spin}_{2n}) = \sum_{U \in P_l^{\pm}(2n)} \Big(\frac{4k^n}{\Pi_k(U)}\Big)^{g-1}.$$

5.5 *Remark.* $P_l(2n)$ may be viewed as a set of highest weights of irreducible representations of Spin_{2n} (namely, those weights in a fundamental chamber for the action of the affine Weyl group of level l). Then P_l^+ is the subset of 'tensor' representations—those which descend to SO_{2n}—and P_l^- is the subset of 'spinor' representations. The same remark applies to the odd spin groups below.

As for SL_4 let us note the lowest cases of this formula. (Note, incidentally, that the formulae (14) are consistent with $\mathrm{Spin}_4 = SL_2 \times SL_2$ and $\mathrm{Spin}_6 = SL_4$: for $n = 2, 3$ respectively N_l coincides with $N_l(SL_2)^2$ and $N_l(SL_4)$.)

$$\begin{array}{lcl} \bullet\ N_1(\mathrm{Spin}_{2n}) & = & 2^{2g} \\ \bullet\ N_2^+(\mathrm{Spin}_{2n}) & = & (2n)^{g-1}(n-1) + 2^{3g-1}n^{g-1} \\ \bullet\ N_2^-(\mathrm{Spin}_{2n}) & = & 2^{3g-1}. \end{array} \tag{14}$$

Proof. We shall just prove the formulae for $N_2^\pm$, and leave N_1 (which we shall not need) to the reader.

We begin by listing the sets over which the summation takes place:

$$\begin{array}{ll} P_2^+(2n): & \{0, 1, \ldots, n-2, n \pm 1\} \\ & \{0, 1, \ldots, n-2, n\} \\ & \cdots \\ & \{0, 2, \ldots, n-1, n\} \\ & \{\pm 1, 2, \ldots, n-1, n\}; \\ & \\ P_2^-(2n): & \{\pm\frac{1}{2}, \frac{3}{2}, \ldots, n \pm \frac{1}{2}\}. \end{array}$$

We recall from [OW] that reflection of the end-points in $0, n$—i.e. the ± signs in these sets—defines an action of $\mathbf{Z}/2 \times \mathbf{Z}/2$ on $P_2(2n)$ under which $\Pi_k(U)$ is easily seen to be invariant. Thus, for example, $P_2^-(2n)$ is a single orbit and so

$$N_2^-(\mathrm{Spin}_{2n}) = 4 \times \Big(\frac{4(2n)^n}{\Pi_{2n}(U)}\Big)^{g-1} \tag{15}$$

where $U = \{\frac{1}{2}, \frac{3}{2} \ldots, n - \frac{1}{2}\}$. We have here

$$\begin{aligned} \Pi_{2n}(U) &= \prod_{1 \le i < j \le n} f_{2n}(i-j) f_{2n}(i+j-1) \\ &= \prod_{r=1}^{2n-1} f_{2n}(r)^{m_r}, \end{aligned}$$

where the multiplicities m_r are to be determined. Namely, $m_r = a_r + b_r$ where a_r is the number of pairs $i < j$ such that $j - i = r$, and b_r is the number of pairs $i < j$ such that $i + j = r + 1$. From this we see that

$$\begin{aligned} a_r &= \begin{cases} n - r & \text{if } r < n \\ 0 & \text{if } r \geq n, \end{cases} \\ b_r &= \begin{cases} [r/2] & \text{if } r < n \\ [(2n - r)/2] & \text{if } r \geq n; \end{cases} \end{aligned}$$

and hence that

$$m_r = n - \Big[\frac{r+1}{2}\Big].$$

So it follows from 5.1 (v) and 5.2 (ii) that

$$\Pi_{2n}(U) = \frac{(2n)^{2n}}{\prod_{r=1}^{2n-1} f(r)^{[\frac{r+1}{2}]}} = \frac{1}{2}(2n)^n;$$

and hence from (15) that $N_2^-(\mathrm{Spin}_{2n}) = 2^{3g-1}$.

Let us now turn to $N_2^+(\mathrm{Spin}_{2n})$. For $l = 0, 1, \ldots, n$ we shall write

$$U_l = \{0, 1, \ldots, \widehat{l}, \ldots, n\},$$

i.e. l has been deleted. Thus $P_2^+(2n)$ consists of $U_0, U_1, \ldots, U_n$ together with the reflections of U_0 and U_n under the action of $\mathbf{Z}/2 \times \mathbf{Z}/2$; and so

$$\begin{aligned} N_2^+(\mathrm{Spin}_{2n}) &= 2\Big(\frac{4(2n)^n}{\Pi_{2n}(U_0)}\Big)^{g-1} + 2\Big(\frac{4(2n)^n}{\Pi_{2n}(U_n)}\Big)^{g-1} \\ &\qquad + \sum_{l=1}^{n-1}\Big(\frac{4(2n)^n}{\Pi_{2n}(U_l)}\Big)^{g-1}. \end{aligned} \tag{16}$$

Now we can write, for each l, $\Pi_{2n}(U_l) = N/D_l$ where:

$$\begin{aligned} N &= \prod_{0 \leq i < j \leq n} f(j-i) f(i+j), \\ D_l &= \prod_{j=l+1}^{n} f(j-l) f(j+l) \prod_{i=0}^{l-1} f(l-i) f(l+i) \quad \text{for } 1 \leq l \leq n-1. \end{aligned}$$

For $l = 0$ and n the denominator takes the slightly simpler forms:

$$D_0 = \prod_{j=1}^{n} f(j)^2 = 16n^2,$$

using 5.1 (iii) and 5.2 (ii); and likewise

$$\begin{aligned} D_n &= \textstyle\prod_{i=0}^{n-1} f(n-i)f(n+i) \\ &= \textstyle\prod_{i=0}^{n-1} f(n-i)^2 \\ &= \textstyle\prod_{i=1}^{n} f(i)^2 \\ &= 16n^2. \end{aligned}$$

To compute the denominator D_l for $1 \leq l \leq n-1$:

$$\begin{aligned} D_l &= f(1)\cdots f(n-l) \times f(2l+1)\cdots f(n+l) \\ &\quad \times f(1)\cdots f(l) \times f(l)\cdots f(2l-1) \\ &= f(n)\frac{f(l)}{f(2l)}\prod_{r=1}^{n-1} f(r)^2 \frac{f(n+1)\cdots f(n+l)}{f(n-1)\cdots f(n-l+1)} \\ &= 4n^2 f(l)f(n+l)/f(2l) && \text{by 5.1 (ii),(iii)} \\ & && \text{and 5.2 (ii);} \\ &= 4n^2 f(n+l)/f(n-l) && \text{using 5.1 (iv);} \\ &= 4n^2. \end{aligned}$$

We next compute the numerator:

$$N = \prod_{r=1}^{2n-1} f(r)^{m_r}$$

where $m_r = a_r + b_r$ and a_r is the number of pairs $i < j$ between 0 and n such that $j - i = r$, and b_r the number such that $i + j = r$. So:

$$\begin{aligned} a_r &= \begin{cases} n+1-r & \text{if } r \leq n \\ 0 & \text{if } r > n, \end{cases} \\ b_r &= \begin{cases} [(r+1)/2] & \text{if } r \leq n \\ [(2n-r+1)/2] & \text{if } r > n; \end{cases} \end{aligned}$$

from which we find

$$m_r = \begin{cases} n+1-[r/2] & \text{if } r \leq n \\ n-[r/2] & \text{if } r > n. \end{cases}$$

Hence

$$N = \prod_{r=1}^{2n-1} f(r)^{n-[r/2]} \times \prod_{r=1}^{n} f(r) = 4(2n)^{n+1},$$

using 5.1 (v) and 5.2 (ii).

Putting together the above computations we obtain $\Pi_{2n}(U_l) = (2n)^{n-1}$ for $l = 0, n$, and $2^{n+1}n^{n-1}$ for $l = 1, \dots, n-1$. Substituting into (16) this gives

$$N_2^+(\mathrm{Spin}_{2n}) = (2n)^{g-1}(n-1) + 2^{3g-1}n^{g-1}$$

as asserted. □

5.6 Example. $\mathbf{Spin_{2n+1}, n \geq 2}$. In this case the Verlinde number is:

$$N_l(\mathrm{Spin}_{2n+1}) = \sum_{U \in P_l(2n+1)} \Big(\frac{4k^n}{\Phi_k(U)}\Big)^{g-1}, \qquad \text{where } k = l + 2n - 1;$$

where $P_l(2n+1)$ consists of $U = \{0 < u_1 < \cdots < u_n\}$ such that

1. $u_i \in \frac{1}{2}\mathbf{Z}$;
2. $u_{i+1} - u_i \in \mathbf{Z}$ for $i = 1, \dots n-1$;
3. $u_{n-1} + u_n < k = l + 2n - 1$.

As in the even case we shall write

$$N_l^{\pm}(\mathrm{Spin}_{2n+1}) = \sum_{U \in P_l^{\pm}} \Big(\frac{4k^n}{\Phi_k(U)}\Big)^{g-1},$$

where $P_l^{\pm}$ denotes the subsets of integral and half-integral U respectively.

Here the lowest Verlinde numbers are:

$$(17) \qquad \begin{array}{lrcl} \bullet & N_1(\mathrm{Spin}_{2n+1}) & = & 2^{g-1}(2^g+1) \\ \bullet & N_2^+(\mathrm{Spin}_{2n+1}) & = & 2^{2g-1} \\ \bullet & N_2^-(\mathrm{Spin}_{2n+1}) & = & (2n+1)^{g-1}(2^{2g-1}+n). \end{array}$$

We shall omit the proof of these formulae, as it is entirely similar to that of (14). Moreover they are proved (by a somewhat clumsier method) in [O2].

5.3 Sections of theta bundles

Our interest in the preceding calculations lies in the fact that $N_2(\mathrm{Spin}_m)$ is the dimension of $H^0(\mathcal{M}(\mathrm{Spin}_m), \Theta(\mathbf{C}^m))$, where $\Theta(\mathbf{C}^m)$ is the theta line bundle, defined in section 2, for the standard orthogonal representation. To each irreducible representation V of a group G there is associated an integer d_V—the *height* or *Dynkin index* of the representation—for which, for any $r \in \mathbf{Z}$, one has

$$h^0(\mathcal{M}(G), \Theta(V)^{\otimes r}) = N_{rd_V}(G). \tag{18}$$

(For a useful discussion of the Dynkin index and the proof of (18) see [LS]; see also [BL], [F], [KNR].)

When $V = \mathbf{C}^m$ is the standard orthogonal representation the height is:

$$d_V = \begin{cases} 4 & \text{if } m = 3, \\ 2 & \text{if } m \geq 5. \end{cases} \tag{19}$$

What is the analogue of (18) for the 'twisted' moduli space $\mathcal{M}^-(\mathrm{Spin}_m)$ defined in section 3? The following is a refinement (which deals with even as well as odd m) of conjecture (5.2) in [OW].

5.7 Conjecture. *For $m \geq 3$, and any representation $SC_m \to SL(V)$ satisfying (6),*

$$h^0(\mathcal{M}^-(\mathrm{Spin}_m), \Theta(V)) = (-1)^m (N^+_{d_V} - N^-_{d_V}).$$

5.8 Example. $\mathbf{m = 3}$. By example 5.3,

$$N_l(\mathrm{Spin}_3) = N_l(SL_2) = \sum_{j=1}^{l+1} \Big(\frac{l+2}{2\sin^2 \frac{j\pi}{l+2}} \Big)^{g-1}.$$

Notice that formally computing 5.6 with $n = 1$ (strictly speaking the Verlinde number 5.6 makes sense only for $n \geq 2$) we find (taking $u_0 = 0$):

$$N_l(\mathrm{Spin}_{2n+1})|_{n=1} = \sum_{j=1}^{2l+1} \Big(\frac{l+1}{4\sin^2 \frac{j\pi}{2l+2}} \Big)^{g-1} = N_{2l}(SL_2).$$

Now, by (18) and the fact that $\mathcal{L}_0 = \Theta(\mathbf{C}^2)$ where $d_{\mathbf{C}^2} = 1$, we have

$$h^0(\mathcal{M}(\mathrm{Spin}_3), \Theta(V)) = h^0(\mathcal{SU}_C(2), \mathcal{L}_0^{d_V}) = N^+_{d_V}(SL_2) + N^-_{d_V}(SL_2).$$

(See example 3.7.)

On the other hand, on $\mathcal{M}^-(\mathrm{Spin}_3) = \mathcal{SU}_C(2,1)$ we have, given a Clifford representation V for which d_V is even, $\Theta(V) = \mathcal{L}_0^{d_V} = \mathcal{L}_1^l$ where $2l = d_V$. Thaddeus's twisted Verlinde formula ([T], corollary (18)) tells us that

$$\begin{aligned} h^0(\mathcal{SU}_C(2,1), \mathcal{L}_1^l) &= \sum_{j=1}^{2l+1} (-1)^{j+1} \Big(\frac{l+1}{4\sin^2 \frac{j\pi}{2l+2}} \Big)^{g-1} \\ &= -N_{2l}^+(SL_2) + N_{2l}^-(SL_2); \end{aligned}$$

or in other words $\mathcal{M}^-(\mathrm{Spin}_3), \Theta(V)) = -N_{d_V}^+(SL_2) + N_{d_V}^-(SL_2)$.

5.9 Example. $\mathbf{m} = \mathbf{6}$. By examples 3.9 and 5.3

$$h^0(\mathcal{M}(\mathrm{Spin}_6), \Theta(V)) = h^0(\mathcal{SU}_C(4), \mathcal{L}_0^{d_V}) = N_{d_V}^+(SL_4) + N_{d_V}^-(SL_4),$$

where

$$h^0(\mathcal{SU}_C(4), \mathcal{L}_0^l) = N_l(SL_4) = \sum_U \Big(\frac{4(l+4)^3}{\Psi_{l+4}(U)} \Big)^{g-1},$$

summed over $U = \{0 = u_0 < u_1 < u_2 < u_3 \le l+3\} \subset \mathbf{Z}$.

On the other hand, $\mathcal{M}^-(\mathrm{Spin}_6) = \mathcal{SU}_C(4,2)$ and by the same arguments as in the previous example, the conjecture is in this case equivalent to:

$$\begin{aligned} h^0(\mathcal{SU}_C(4,2), \mathcal{L}_2^l) &= N_{2l}^+(SL_4) - N_{2l}^-(SL_4) \\ &= \sum_U (-1)^{u_2} \Big(\frac{4(2l+4)^3}{\Psi_{2l+4}(U)} \Big)^{g-1}, \end{aligned}$$

summed over $U = \{0 = u_0 < u_1 < u_2 < u_3 \le 2l+3\} \subset \mathbf{Z}$. One can possibly verify this using [BL] theorem 9.4.

6 Numerology

Our principal aim in this section to make some sense of the Verlinde numbers (13), (14) and (17) computed in the previous section at level 2—those, that is, associated to the orthogonal representation of the Clifford group. But we begin with a remark, independent of the rest of the paper, which deals with the Verlinde number (17) at level 1.

6.1 A remark on the generator of Pic $\mathcal{M}(\mathrm{Spin}_{2n+1})$

It is known from [LS] that the Picard group of the moduli scheme $\mathcal{M}(\mathrm{Spin}_m)$ is infinite cyclic. When $m = 2n+1$ is odd the ample generator $\mathcal{P}$ is constructed as a Pfaffian bundle, i.e. $2\mathcal{P} = \Theta(\mathbf{C}^{2n+1})$. (When $m = 2n$ is even $\mathcal{P}$ exists only as a Weil divisor class.) Then according to (17) the space of sections of this line bundle has dimension

$$h^0(\mathcal{M}(\mathrm{Spin}_{2n+1}), \mathcal{P}) = N_1(\mathrm{Spin}_{2n+1}) = 2^{g-1}(2^g+1).$$

This formula is striking, first because it is independent of n and second because it is the number of even theta characteristics of the curve. In fact it is easy to see how to construct a basis for the linear system $|\mathcal{P}|$, as follows.

Let $\vartheta^+(C)$ denote the set of even theta characteristics, and for each $L \in \vartheta^+(C)$ consider the reduced divisor

$$D_L = \{E \in \mathcal{M}(\mathrm{Spin}_{2n+1}) | H^0(C, L \otimes E(\mathbf{C}^{2n+1})) \neq 0\}.$$

By proposition 4.6 the dimension of $H^0(C, L \otimes E(\mathbf{C}^{2n+1}))$ is always even, and so by theorem 4.12 D_L is a proper subset of the moduli space. Consequently it is divisor—this is shown in proposition 7.1 below—and in fact, since we take D_L to be reduced, some multiple $kD_L \in |\Theta(\mathbf{C}^{2n+1})| = |2\mathcal{P}|$. Since the dimension of $H^0(C, L \otimes E(\mathbf{C}^{2n+1}))$ is always even we see that k is at least, and hence equal to, 2. We have therefore constructed a set of divisors on $\mathcal{M}(\mathrm{Spin}_{2n+1})$:

$$D_L \in |\mathcal{P}|, \qquad L \in \vartheta^+(C).$$

In the case $n = 1$ these were shown by Beauville [B2] to be linearly independent, and hence a basis of the linear system; we expect the same to be true in general.

6.2 Prym varieties

We recall the following ‘Verlinde numbers’ for principally polarised abelian varieties. Let (A, Ξ) be any principally polarised abelian variety of dimension g, where Ξ is a symmetric divisor representing the polarisation; and let

$$H^0(A, m\Xi) = H^0_+(A, m\Xi) \oplus H^0_-(A, m\Xi)$$

be the decomposition into ±-eigenspaces under the canonical involution of A. Then by writing down a suitable basis of theta functions one can easily verify that:

$$\dim H^0_\pm(A, m\Xi) = \begin{cases} (m^g \pm 2^g)/2 & \text{if } m \equiv 0 \bmod 2, \\ (m^g \pm 1)/2 & \text{if } m \equiv 1 \bmod 2. \end{cases} \tag{20}$$

Associated to a smooth projective curve C we have a natural configuration of principally polarised *Prym varieties.* Let us recall the usual notation (see [ACGH]). For each nonzero half-period $\eta \in J_2(C)\backslash\{\mathcal{O}\}$ we have an unramified double cover

$$p : \widetilde{C}_\eta \to C.$$

Writing $\widetilde{J}^{2g-2} = J^{2g-2}(\widetilde{C}_\eta)$ we have

$$\mathrm{Nm}_p^{-1}(K_C) = P_\eta \cup P_\eta^- \subset \widetilde{J}^{2g-2};$$

where P_η, P_η^- are disjoint translates of the same abelian subvariety, characterised by the condition that for $L \in \mathrm{Nm}_p^{-1}(K_C)$:

$$h^0(\widetilde{C}_\eta, L) \equiv \begin{cases} 0 & \bmod 2 \text{ if } L \in P_\eta, \\ 1 & \bmod 2 \text{ if } L \in P_\eta^-. \end{cases} \tag{21}$$

Then P_η is called the Prym variety of the covering. We shall denote by Ξ_η the symmetric divisor representing the canonical principal polarisation on P_η, defined by $2\Xi_\eta = P_\eta \cap \widetilde{\Theta}$, where $\widetilde{\Theta}$ is the theta-divisor in $\widetilde{J}^{2g-2}$. We shall allow also $\eta = 0$ by setting $(P_0, \Xi_0) = (J^{g-1}(C), \theta)$.

Notice that on P_η^- both the polarisation and the line bundle $2\Xi_\eta$ are defined, and identify under translation with the same objects on P_η. The distinguished line bundle Ξ_η which represents the polarisation is no longer defined on P_η^-, however. Thus in what follows the line bundle $m\Xi_\eta$ makes sense on P_η^- *only if m is even.*

6.3 Spin Verlinde numbers versus Prym Verlinde numbers

We may now formulate the computations of section 5 as follows.

6.1 Theorem. *1. If $m \geq 6$ is even then*

$$\begin{aligned} N_2(\mathrm{Spin}_m) = N_2^+ + N_2^- &= \sum_{\eta \in J_2(C)} h^0_+(P_\eta, m\Xi_\eta) + \sum_{\eta \neq 0} h^0_+(P_\eta^-, m\Xi_\eta), \\ N_2^+ - N_2^- &= \sum_{\eta \in J_2(C)} h^0_-(P_\eta, m\Xi_\eta) + \sum_{\eta \neq 0} h^0_-(P_\eta^-, m\Xi_\eta). \end{aligned}$$

2. If $m \geq 5$ is odd then

$$\begin{aligned} N_2(\mathrm{Spin}_m) = N_2^+ + N_2^- &= \sum_{\eta \in J_2(C)} h^0_+(P_\eta, m\Xi_\eta), \\ -N_2^+ + N_2^- &= \sum_{\eta \in J_2(C)} h^0_-(P_\eta, m\Xi_\eta). \end{aligned}$$

Proof. Equations (13), (14), (17) and (20). □

6.2 *Remarks. (i)* Of course, when m is even P_η and P_η^- are interchangeable as far as the dimensions alone are concerned. But in view of the constructions which follow in the next sections, together with the remarks in section 7.1, it will be seen that the right-hand side of part 1 is the correct way to write these identities.

(ii) For $m = 3$ there is an analogous statement obtained by replacing N_2 by N_4. This case has been discussed at length in [OP].

These results imply, via (18), (19) and 5.7 (assuming the validity of the latter) identities among h^0s which we can summarise in the following table.

(22)

	$h^0(\mathcal{M}(\mathrm{Spin}_m), \Theta(\mathbf{C}^m))$	$h^0(\mathcal{M}^-(\mathrm{Spin}_m), \Theta(\mathbf{C}^m))$
odd m	$\sum_{\eta \in J_2} h^0_+(P_\eta, m\Xi_\eta)$	$\sum_{\eta \in J_2} h^0_-(P_\eta, m\Xi_\eta)$
even m	$\sum_{\eta \in J_2} h^0_+(P_\eta, m\Xi_\eta)$ $+ \sum_{\eta \in J_2 \setminus \{0\}} h^0_+(P_\eta^-, m\Xi_\eta)$	$\sum_{\eta \in J_2} h^0_-(P_\eta, m\Xi_\eta)$ $+ \sum_{\eta \in J_2 \setminus \{0\}} h^0_-(P_\eta^-, m\Xi_\eta)$

Although theorem 6.1 is stated only for $m \geq 5$, table (22) is in fact valid for *all* $m \in \mathbf{N}$, as we shall see next by a case-by-case examination.

6.3 Example. m = 1. We can identify $\mathcal{M}(\mathrm{Spin}_1)$ with $J_2(C)$; while $\mathcal{M}^-(\mathrm{Spin}_1)$ is empty. So (22) is trivial in this case.

6.4 Example. m = 2. From example 3.6, $\mathcal{M}^{\pm}(\mathrm{Spin}_2)$ are both copies of $\mathrm{Pic}(C)$. However, the orthogonal theta bundle $\Theta(\mathbf{C}^2)$ is defined *only* on the degree 0 component $J(C)$ of $\mathcal{M}(\mathrm{Spin}_2)$, where it can be identified with the line bundle 8θ where θ is a theta divisor on the Jacobian. So in this case again one may readily check from (20) that the identity in (22) holds. But in fact we have more:

$$\begin{aligned} H^0_+(J(C), 8\theta) &\cong \sum_{\eta\in J_2} H^0(P_\eta, 2\Xi_\eta), \\ H^0_-(J(C), 8\theta) &\cong \sum_{\eta\in J_2\backslash\{0\}} H^0(P^-_\eta, 2\Xi_\eta). \end{aligned}$$

One can see this as follows. Start with the fact that for any symmetric line bundle $\mathcal{L}$ on an abelian variety A one has

$$H^0(A, \mathcal{L}^4) \cong \bigoplus_{\alpha\in\widehat{A}_2} H^0(A, \mathcal{L}\otimes\alpha)$$

where $\widehat{A}_2 = \{\alpha \in \mathrm{Pic}^0 A | \alpha^2 = \mathcal{O}_A\}$. This isomorphism is obtained by pulling back sections under the squaring map $[2] : A \to A$; for each α we have $[2]^*(\mathcal{L}\otimes\alpha) = \mathcal{L}^4$. Noting that $[2]$ commutes with the involution $[-1] : A \leftrightarrow A$ we see that in fact $H^0_\pm(A, \mathcal{L}^4) \cong \bigoplus_{\alpha\in\widehat{A}_2} H^0_\pm(A, \mathcal{L}\otimes\alpha)$.

If θ represents a principal polarisation on A then we can identify $A_2 \cong \widehat{A}_2$ by $\eta \mapsto \theta_\eta - \theta$ where $\theta_\eta = t^*_\eta\theta$. In particular, when $A = J(C)$ and $\mathcal{L} = 2\theta$ we obtain

$$H^0_\pm(J, 8\theta) \cong \bigoplus_{\eta\in J_2} H^0_\pm(J, \theta+\theta_\eta).$$

On the other hand $H^0_\pm(J, \theta+\theta_\eta) \cong H^0(P^\pm_\eta, 2\Xi_\eta)$ for $\eta \neq 0$. To see this choose $\zeta \in J(C)$ such that $\zeta^2 = \eta$. Then translation by ζ induces an isomorphism $t^*_\zeta : H^0(J, \theta+\theta_\eta) \xrightarrow{\sim} H^0(J, 2\theta)$, which is equivariant with respect to $[-1]^*$ acting on the first space and t^*_η acting on the second. Hence the spaces $H^0_\pm(J, \theta+\theta_\eta)$ identify with the $\pm$-eigenspaces in $H^0(J, 2\theta)$ under t^*_η; and these in turn are classically identified with $H^0(P^\pm_\eta, 2\Xi_\eta)$ by the Schottky-Jung relations.

6.5 Example. $\mathbf{m = 3}$. We have seen (see example 3.7) that $\Theta(\mathbf{C}^3)$ restricts to $\mathcal{L}_0^4 \to \mathcal{SU}_C(2) = \mathcal{M}(\mathrm{Spin}_3)$ and to $\mathcal{L}_1^2 \to \mathcal{SU}_C(2,1) = \mathcal{M}^-(\mathrm{Spin}_3)$—in each case the anticanonical line bundle. The coincidence of dimensions in (22) for this case was the first to be observed, and has been explored in [OP] and [R2].

6.6 Example. $\mathbf{m = 4}$. By example 3.8 we have $\mathcal{M}(\mathrm{Spin}_4) \cong \mathcal{SU}_C(2) \times \mathcal{SU}_C(2)$; while the 4-dimensional orthogonal representation is the tensor product of the 2-dimensional representations of the distinct factors, from which the theta bundle in (22) is $\Theta(\mathbf{C}^4) = pr_+^* \mathcal{L}_0^2 \otimes pr_-^* \mathcal{L}_0^2$, where $\mathcal{L}_0 = \Theta(\mathbf{C}^2)$ is the ample generator of Pic $\mathcal{SU}_C(2) = \mathbf{Z}$. Thus

$$H^0(\mathcal{M}(\mathrm{Spin}_4), \Theta(\mathbf{C}^4)) \cong \bigotimes^2 H^0(\mathcal{SU}_C(2), \mathcal{L}_0^2),$$

and since $h^0(\mathcal{SU}_C(2), \mathcal{L}_0^2) = 2^{g-1}(2^g + 1)$ we can at once verify the case $m = 4$ of the table. Moreover, just as for the case $m = 2$ we can say more:

$$\begin{aligned} S^2 H^0(\mathcal{SU}_C(2), \mathcal{L}_0^2) &\cong \sum_{\eta \in J_2} H^0_+(P_\eta, 4\Xi_\eta), \\ \bigwedge^2 H^0(\mathcal{SU}_C(2), \mathcal{L}_0^2) &\cong \sum_{\eta \in J_2 \setminus \{0\}} H^0_+(P_\eta^-, 4\Xi_\eta). \end{aligned}$$

As in the case $m = 2$ we have written down not just an identity of dimensions, but in fact an isomorphism of vector spaces. In this case it follows easily (for C without vanishing theta-nulls) from the results of [B2].

Similarly $\mathcal{M}^-(\mathrm{Spin}_4)$ is isomorphic to $\mathcal{SU}_C(2,1) \times \mathcal{SU}_C(2,1)$, with theta bundle $\Theta(\mathbf{C}^4) \cong pr_+^* \mathcal{L}_1 \otimes pr_-^* \mathcal{L}_1$, where $\mathcal{L}_1$ is the generator of Pic $\mathcal{SU}_C(2,1)$. This time $h^0(\mathcal{SU}_C(2,1), \mathcal{L}_1^2) = 2^{g-1}(2^g - 1)$, the results of Beauville can be applied to give isomorphisms

$$\begin{aligned} S^2 H^0(\mathcal{SU}_C(2,1), \mathcal{L}_1) &\cong \sum_{\eta \in J_2} H^0_-(P_\eta, 4\Xi_\eta), \\ \bigwedge^2 H^0(\mathcal{SU}_C(2,1), \mathcal{L}_1) &\cong \sum_{\eta \in J_2 \setminus \{0\}} H^0_-(P_\eta^-, 4\Xi_\eta), \end{aligned}$$

and again table (22) is verified in this case.

6.7 Example. m = 6. In this case example 3.9 identifies $\Theta(\mathbf{C}^6) \to \mathcal{M}^{\pm}(\mathrm{Spin}_6)$ with $\mathcal{L}_0^2 \to \mathcal{SU}_C(4)$ and $\mathcal{L}_2 \to \mathcal{SU}_C(4,2)$ respectively; and table (22) for the case $\mathcal{M}^+(\mathrm{Spin}_6)$ suggests isomorphisms

$$\begin{array}{rcl} H^0_+(\mathcal{SU}_C(4),\mathcal{L}_0^2) & \cong & \sum_{\eta\in J_2} H^0_+(P_\eta, 6\Xi_\eta), \\ H^0_-(\mathcal{SU}_C(4),\mathcal{L}_0^2) & \cong & \sum_{\eta\in J_2\backslash\{0\}} H^0_+(P_\eta^-, 6\Xi_\eta); \end{array}$$

for *some* ±-decomposition of $H^0(\mathcal{SU}_C(4),\mathcal{L}_0^2)$. We shall observe in remark 7.1 in the next section that the involution of $\mathcal{M}(\mathrm{Spin}_6)$ analogous to the exchange of factors in example 6.6 is the dualising involution $E \mapsto E^\vee$ of $\mathcal{SU}_C(4)$. We therefore conjecture isomorphisms as above where $H^0_\pm$ on the left-hand side are the ±-eigenspaces for the dualising involution.

For $\mathcal{M}^-(\mathrm{Spin}_6) = \mathcal{SU}_C(4,2)$, remark 7.1 will say that the analogous involution is $E \mapsto E^\vee \otimes \mathrm{Nm}E$, where $\mathrm{Nm}E$ is the fixed spinor norm, satisfying $(\mathrm{Nm}E)^2 = \det E$. Then we expect, in this case:

$$\begin{array}{rcl} H^0_+(\mathcal{SU}_C(4,2),\mathcal{L}_2) & \cong & \sum_{\eta\in J_2} H^0_-(P_\eta, 6\Xi_\eta), \\ H^0_-(\mathcal{SU}_C(4,2),\mathcal{L}_2) & \cong & \sum_{\eta\in J_2\backslash\{0\}} H^0_-(P_\eta^-, 6\Xi_\eta). \end{array}$$

7 Constructing the homomorphisms: the Jacobian

Our aim in this section and the next is to construct homomorphisms:

$$H^0(\mathcal{M}^{\pm}(\mathrm{Spin}_{2n+1}),\Theta(\mathbf{C}^{2n+1}))^\vee \to \sum_{\eta\in J_2} H^0_\pm(P_\eta,(2n+1)\Xi_\eta) \tag{23}$$

$$\begin{array}{rcl} H^0(\mathcal{M}^{\pm}(\mathrm{Spin}_{2n}),\Theta(\mathbf{C}^{2n}))^\vee & \to & \sum_{\eta\in J_2} H^0_\pm(P_\eta, 2n\Xi_\eta) \\ & & \oplus \sum_{\eta\neq 0} H^0_\pm(P_\eta^-, 2n\Xi_\eta) \end{array} \tag{24}$$

Table (22) says that in each case the left- and right-hand sides have the same dimension, and so we naturally conjecture that these homomorphisms are isomorphisms, though we shall not prove this here. As remarked in the last section, the first few cases $m = 1,2,3,4$ are well understood, and we do have natural isomorphisms (23), (24) in these cases.

7.1 The splitting for even spin groups

We shall at this point attempt to explain the splitting $H^0(\Theta(\mathbf{C}^m)) = \sum_{\eta \in J_2} \oplus \sum_{\eta \neq 0}$ when m is even. Note that for even m the group Spin_m carries a unique nontrivial outer automorphism:



This induces an involution $\sigma : \mathcal{M}(SC_m) \leftrightarrow \mathcal{M}(SC_m)$ which preserves the spinor norm and hence acts on $\mathcal{M}(\mathrm{Spin}_m)$ and $\mathcal{M}^-(\mathrm{Spin}_m)$. (More concretely, the group automorphism is obtained by Clifford conjugation by a vector in $\mathbf{C}^m$; when m is *odd* this is an inner automorphism. σ then acts on Clifford bundles by conjugating transition functions with respect to some open cover; when m is odd this action is still defined, of course, but is trivial.)

On the other hand this automorphim preserves the orthogonal representation, and hence the isomorphism class of the line bundle $\Theta(\mathbf{C}^m)$; accordingly $H^0(\Theta(\mathbf{C}^m)$ splits into $\pm$-eigenspaces $H^0_+ \oplus H^0_-$ under the action of σ.

For even m we now expect the following refinement of (24):

$$\begin{aligned} H^0_+(\mathcal{M}^{\pm}(\mathrm{Spin}_m), \Theta(\mathbf{C}^m))^{\vee} &\rightarrow \sum_{\eta \in J_2} H^0_{\pm}(P_\eta, m\Xi_\eta), \\ H^0_-(\mathcal{M}^{\pm}(\mathrm{Spin}_m), \Theta(\mathbf{C}^m))^{\vee} &\rightarrow \sum_{\eta \neq 0} H^0_{\pm}(P^-_\eta, m\Xi_\eta). \end{aligned} \tag{25}$$

Note that this is exactly what happens in the case $m = 4$ (see example 6.6), where σ simply exchanges factors in the products $\mathcal{M}(\mathrm{Spin}_4) = \mathcal{SU}_C(2) \times \mathcal{SU}_C(2)$ and $\mathcal{M}^-(\mathrm{Spin}_4) = \mathcal{SU}_C(2,1) \times \mathcal{SU}_C(2,1)$, so that the summands $H^0_{\pm}$ in (25) are precisely $S^2 H^0$ and $\bigwedge^2 H^0$ appearing in 6.6.

In the case $m = 6$—recall example 3.9—we have $\mathcal{M}(\mathrm{Spin}_6) = \mathcal{SU}_C(4)$ and $\mathcal{M}^-(\mathrm{Spin}_6) = \mathcal{SU}_C(4,2)$. In each case not only the determinant $\det E$ of vector bundles E is fixed, but also a square root $\mathrm{Nm} E$ of $\det E$. Rank 4 vector bundles come from SC_6-bundles via the first half-spinor representation $SC_6 \rightarrow GL_4$; one easily checks that taking instead the second projection to GL_4 induces the involution on rank 4 vector bundles $\sigma : E \leftrightarrow E^{\vee} \otimes \mathrm{Nm} E$.

Thus in the case $m = 6$ of (25) we expect the situation already described in example 6.7 of the previous section.

7.2 The main construction

We shall construct the homomorphisms (23), (24) one summand at a time, concentrating in this section on the summand $\eta = 0$, i.e. the projection to Jacobian thetas (see (27) and corollary 7.4). In the next section section we shall construct the remaining projections to the Prym thetas (corollary 8.7).

It will be convenient to denote the two-component variety $\mathcal{M}(\mathrm{Spin}_m) \cup \mathcal{M}^-(\mathrm{Spin}_m)$ by $\mathcal{N}(m)$. Let $\rho : SC_m \to SL(V)$ be a representation satisfying condition (6) from section 2. (We are *mainly* concerned with the orthogonal representation $V = \mathbf{C}^m$, and $r = m$ in proposition 7.1 below.)

We consider the subset

$$\mathcal{D}(V) = \{(L, E) | H^0(C, L \otimes E(V)) \neq 0\} \subset J^{g-1}(C) \times \mathcal{N}(m). \tag{26}$$

It is most important for us that in the case $V = \mathbf{C}^m$ is the vector representation—by theorem 4.10—this is a proper subset in each component of the product.

For the theta divisor θ in $J^{g-1}(C)$ and the theta bundle $\Theta(V)$ on $\mathcal{N}(m)$ we shall abuse notation and use the same symbols to denote also their pull-back to the product $J^{g-1}(C) \times \mathcal{N}(m)$.

7.1 Proposition. *If $H^0(C, L \otimes E(V))$ is generically zero—as it is when $V = \mathbf{C}^m$—then $\mathcal{D}(V)$ is the support of a divisor in $|r\theta + \Theta(V)|$, where $r = \dim V$.*

Proof. The representation ρ induces a morphism of varieties

$$\alpha : \mathcal{N}(m) \to \mathcal{M}(SC_m) \to \mathcal{M}(SL_r);$$

and by definition $\mathcal{D}(V)$ is (the support of) the pull-back via

$$J^{g-1}(C) \times \mathcal{N}(m) \xrightarrow{id \times \alpha} J^{g-1}(C) \times \mathcal{M}(SL_r) \xrightarrow{\otimes} \mathcal{U}(r, r(g-1))$$

(where, of course, we are identifying $\mathcal{M}(SL_r)$ with the moduli space of semistable vector bundles of rank r and trivial determinant, and the second map is tensor product of vector bundles) of the canonical theta divisor (see section 2)

$$\Theta_{r,r(g-1)} = \{F | H^0(F) \neq 0\} \subset \mathcal{U}(r, r(g-1)).$$

Since by hypothesis $H^0(C, L \otimes E(V)) = 0$ generically, it follows that $\mathcal{D}(V)$ is a well-defined divisor. The proposition now follows from the discussion of section 2: first of all, the pull-back of $\Theta_{r,r(g-1)}$ to $J^{g-1}(C) \times \mathcal{M}(SL_r)$ restricts on a fibre $\{L\} \times \mathcal{M}(SL_r)$ as the restriction of $\Theta_{r,0}$ from $\mathcal{U}(r, 0)$—this is by definition of $\Theta_{r,0}$—and we have seen that this is just $\Theta(\mathbf{C}^r)$. Hence the pull-back to $J^{g-1}(C) \times \mathcal{N}(m)$ is $\Theta(V)$ on fibres $\{L\} \times \mathcal{N}(m)$.

On the other hand, restriction to fibres $J^{g-1}(C) \times \{V\}$, for any vector bundle V with rank r and trivial determinant, is well-known to be independent of V—see for example [OP] section 3.1. Then it follows from (5) that $\Theta_{r,r(g-1)}$ restricts to $r\theta$. $\square$

It follows from the Künneth theorem that $\mathcal{D}(V)$ defines up to scalar a tensor in

$$H^0(\mathcal{N}(m), \Theta(V)) \otimes H^0(J^{g-1}, r\theta),$$

or equivalently a homomorphism, the projection of (23), (24) at $\eta = 0$,

$$s_0 : H^0(\mathcal{N}(m), \Theta(V))^\vee \to H^0(J^{g-1}, r\theta). \tag{27}$$

s_0 is dual to pull-back of hyperplane sections under the rational map $f_0 : \mathcal{N}(m) \to |r\theta|$ sending E to $\mathcal{D}(V)|_{J^{g-1}\times\{E\}}$.

Now suppose that the vector bundles $E(V)$ are self-dual. This happens when the representation V is symplectic or orthogonal—in particular if $V = \mathbf{C}^m$ or a spin representation. Then it follows easily from Riemann-Roch and Serre duality that the divisors $f_0(E)$ are *symmetric*:

$$f_0 : \mathcal{N}(m) \to |r\theta|_+ \cup |r\theta|_-$$

where $|r\theta|_\pm = \mathbf{P}H^0_\pm(J^{g-1}, r\theta)$.

7.2 Proposition. *When $V = \mathbf{C}^m$ is the standard orthogonal representation, for $m \geq 3$, f_0 respects parity: $f_0 : \mathcal{M}^\pm(\mathrm{Spin}_m) \to |m\theta|_\pm$ respectively.*

Before proving this, we need to make a general remark about principally polarised abelian varieties (A, Ξ), where as usual Ξ is a symmetric divisor representing the polarisation. Let $\vartheta(A) = A_2$ denote the set of 2-torsion points, and let

$$\begin{aligned} \vartheta^+(A) &= \{x \in A_2 \mid \mathrm{mult}_x \Xi \equiv 0 \bmod 2\} \\ \vartheta^-(A) &= \{x \in A_2 \mid \mathrm{mult}_x \Xi \equiv 1 \bmod 2\} \end{aligned}$$

In the case $(A, \Xi) = (J^{g-1}, \theta)$ we shall write $\vartheta^{\pm}(J^{g-1}) = \vartheta^{\pm}(C)$. These are the sets of even and odd theta characteristics.

In the case of a Prym variety (P_η, Ξ_η) it is shown in [OP], proposition 2.3 that we can identify:

$$\begin{aligned} \vartheta^{\pm}(P_\eta) \;&=\; \{\pi^* N = \pi^*(\eta \otimes N) \in \widetilde{J}^{2g-2} \\ &\qquad \text{where } N, \eta \otimes N \in \vartheta^{\pm}(C)\}. \end{aligned} \tag{28}$$

On P_η^- there is again an induced principal polarisation, though, as remarked in section 6.2, no distinguished theta divisor. Thus we may talk about $\vartheta(P_\eta^-)$, whose points are described (using the same methods as in [OP]) by:

$$\begin{aligned} \vartheta(P_\eta^-) \;&=\; \{\pi^* N = \pi^*(\eta \otimes N) \in \widetilde{J}^{2g-2} \\ &\qquad \text{where } N, \eta \otimes N \in \vartheta(C) \text{ have } \textit{opposite} \text{ parity}\}. \end{aligned} \tag{29}$$

But for P_η^- the partition into $\vartheta^{\pm}$ is no longer well-defined.

7.3 Lemma. *1. If m is odd then $|m\Xi|_+$ (resp. $|m\Xi|_-$) is the linear subsystem with base-point set $\vartheta^-(A)$ (resp. $\vartheta^+(A)$).*

2. If m is even then $|m\Xi|_+$ is base-point free; while $|m\Xi|_-$ is the linear subsystem with base-point set A_2.

Proof. See [OP] lemma 2.2 (for part 1); or [LB] chapter 4 section 7. □

Proof of proposition 7.2. We use proposition 4.6. Suppose first that m is odd: then for $E \in \mathcal{N}(m)$ and any theta characteristic $L \in \vartheta(C)$ we have

$$h^0(C, L \otimes E(\mathbf{C}^m)) \equiv h^0(C, L) + \deg \mathrm{Nm} E \mod 2.$$

So if $E \in \mathcal{M}(\mathrm{Spin}_m)$ then by definition $L \in f_0(E)$ for all *odd* theta characteristics; so by the first part of the lemma $f_0(E) \in |m\theta|_+$. Likewise $f_0(E) \in |m\theta|_-$ whenever $E \in \mathcal{M}^-(\mathrm{Spin}_m)$.

If m is even then the same argument shows that $\mathcal{M}^-(\mathrm{Spin}_m)$ maps into $|m\theta|_-$. $\mathcal{M}(\mathrm{Spin}_m)$, on the other hand, maps either into $|m\theta|_+$ or $|m\theta|_-$, and to see that it is not the latter it suffices to exhibit a theta characteristic $K^{\frac{1}{2}}$ and bundle $E \in \mathcal{M}(\mathrm{Spin}_m)$ for which $H^0(C, K^{\frac{1}{2}} \otimes E(\mathbf{C}^m)) = 0$—which is possible by theorem 4.12. □

7.4 Corollary. *When $V = \mathbf{C}^m$ is the standard orthogonal representation the homomorphism s_0 respects parity:*

$$s_0 : H^0(\mathcal{M}^{\pm}(\mathrm{Spin}_m), \Theta(\mathbf{C}^m))^\vee \to H^0_{\pm}(J^{g-1}, m\theta) \qquad \textit{respectively}.$$

8 Constructing the homomorphisms: the Pryms

We next explain how the construction of the previous section may be extended to give the projections to the remaining summands, $\eta \neq 0$, in (23) and (24). These are given in corollary 8.7, which extends corollary 7.4.

To begin, it is necessary to note that semistability of a bundle is preserved under pull-back to the double covers.

8.1 Lemma. *Let $p : \widetilde{C} \to C$ be any unramified cover of smooth projective curves. Then, if a vector bundle $V \to C$ is semistable then $p^*V \to \widetilde{C}$ is semistable.*

Proof. By the Narasimhan-Seshadri theorem V is induced from a projective unitary representation of the fundamental group $\pi_1(C)$. Since $\widetilde{C}$ is an unramified cover its fundamental group injects into $\pi_1(C)$, and the restriction of the above representation then induces the pull-back bundle, which is consequently semistable. □

8.2 Corollary. *Let $p : \widetilde{C} \to C$ be as in the previous lemma, and $E \to C$ a semistable G-bundle. Then $p^*E \to \widetilde{C}$ is semistable.*

Proof. The same argument as in the above proof works for G-bundles by Ramanathan's generalisation of the Narasimhan-Seshadri theorem [Rth1]; alternatively apply the lemma to the adjoint bundle ad E: by [Rth2], corollary 2.18, semistability of E is equivalent to semistability of ad E as a vector bundle. □

Let us now return to the double cover $p : \widetilde{C}_\eta \to C$. Noting that for a Clifford bundle $E \to C$ the spinor norm satisfies $\mathrm{Nm}(p^*E) = p^*\mathrm{Nm}(E)$, and this has even degree, it follows from corollary 8.2 that we obtain a morphism of moduli spaces

$$u = p^* : \mathcal{N}_C(m) \to \mathcal{M}_{\widetilde{C}_\eta}(\mathrm{Spin}_m).$$

8.3 Proposition. *For any representation* $SC_m \to SL(V)$ *we have* $u^*\Theta_{\mathcal{M}_{\widetilde{C}_\eta}}(V) = 2\Theta_{\mathcal{N}_C}(V)$.

Proof. Let $E \to C \times S$ be an arbitrary family of semistable SC_m-bundles, and let $F = E(V)$ be the associated family of vector bundles via the given representation. Let $\widetilde{F} = (p \times \mathrm{id})^* F$ be the pull-back of the family by the double cover:

$$\begin{tikzcd}
\widetilde{C}_\eta \times S \arrow[rr, "p\times \mathrm{id}"] \arrow[dr, "\widetilde{\pi}"'] & & C \times S \arrow[dl, "\pi"] \\
& S. &
\end{tikzcd}$$

It is clear from the discussion of section 2 that to prove the proposition it suffices to show that

$$\Theta(\widetilde{F}) = 2\Theta(F) :$$

i.e. the line bundle $\Theta(V) \to \mathcal{M}_C(SC_m)$ represents the functor $E \mapsto \Theta(F)$, while the line bundle $u^*\Theta_{\mathcal{M}_{\widetilde{C}_\eta}}(V) \to \mathcal{M}_C(SC_m)$ represents the functor $E \mapsto \Theta(\widetilde{F})$.

So to compute $\Theta(\widetilde{F})$, first note that by the projection formula applied to $p \times \mathrm{id}$ we have, for $i = 0, 1$:

$$\begin{aligned}
R^i_{\widetilde{\pi}}(\widetilde{F}) &= R^i_\pi(F \otimes p_*\mathcal{O}_{\widetilde{C}_\eta}) \\
&= R^i_\pi(F \oplus F \otimes \eta) \\
&= R^i_\pi(F) \oplus R^i_\pi(F \otimes \eta).
\end{aligned}$$

If we fit the direct images $R^i_\pi(F)$ into an exact sequence (1), and $R^i_\pi(F \otimes \eta)$ into a similar sequence with middle terms $K^{0\prime} \xrightarrow{\phi'} K^{1\prime}$, then we get an exact sequence:

$$0 \to R^0_{\widetilde{\pi}}\widetilde{F} \to K^0 \oplus K^{0\prime} \xrightarrow{\mathrm{diag}(\phi,\phi')} K^1 \oplus K^{1\prime} \to R^1_{\widetilde{\pi}}\widetilde{F} \to 0.$$

It follows at once that

$$\mathrm{Det}(\widetilde{F}) = \mathrm{Det}(F) \otimes \mathrm{Det}(F \otimes \eta).$$

But since the bundle F has trivial determinant we can replace Det by Θ here. And since $\Theta(F \otimes \eta) = \Theta(F)$ by corollary 2.4, we obtain $\Theta(\widetilde{F}) = 2\Theta(F)$ as required. □

We now consider

$$\widetilde{\mathcal{D}}(V) \subset \widetilde{J}^{2g-2} \times \mathcal{M}_{\widetilde{C}_\eta}(\mathrm{Spin}_m),$$

denoting the divisor of (26) and proposition 7.1 with C replaced by $\widetilde{C}_\eta$. As a consequence of proposition 8.3, we see that the pull-back via the map

$$\mathrm{incl} \times u : (P_\eta \cup P_\eta^-) \times \mathcal{N}(m) \to \widetilde{J}^{2g-2} \times \mathcal{M}_{\widetilde{C}_\eta}(\mathrm{Spin}_m),$$

of $\widetilde{\mathcal{D}}(V)$ is—*provided* $H^0(\widetilde{C}_\eta, L \otimes p^*E(V)) = 0$ *for generic* $(L, E) \in P_\eta^\pm \times \mathcal{M}^\pm(\mathrm{Spin}_m)$—a divisor

$$\mathcal{E}_\eta(V) \in |2r\Xi_\eta + 2\Theta_\mathcal{N}|. \tag{30}$$

In the case of the orthogonal representation—except possibly for $\mathcal{M}^-(\mathrm{Spin}_{2n})$—this is guaranteed by corollary 4.13:

8.4 Proposition. *Let* $V = \mathbf{C}^m$ *be the standard orthogonal representation.*

1. *If* m *is odd and* $L \in P_\eta^-$ *then* $H^0(\widetilde{C}_\eta, L \otimes p^*E(V)) \neq 0$ *for all* $E \in \mathcal{M}^\pm(\mathrm{Spin}_m)$.
2. *In all other cases—except possibly* $E \in \mathcal{M}^-(\mathrm{Spin}_{2n})$*—we have* $H^0(\widetilde{C}_\eta, L \otimes p^*E(V)) = 0$ *for generic* $(L, E) \in P_\eta \times \mathcal{M}^\pm(\mathrm{Spin}_m)$ *or* $P_\eta^- \times \mathcal{M}^\pm(\mathrm{Spin}_m)$.

Proof. Note that by corollary 4.7 (applied to $\widetilde{C}_\eta$), together with (21), $h^0(\widetilde{C}_\eta, L \otimes p^*E(\mathbf{C}^m))$ is odd if m is odd and $L \in P_\eta^-$, hence nonzero as asserted. In all other cases, on the other hand, $h^0(\widetilde{C}_\eta, L \otimes p^*E(\mathbf{C}^m))$ is even. So the proposition is equivalent to 4.13. □

The next step is to observe that the divisor thus constructed has multiplicity two:

8.5 Proposition. *Suppose that the representation* V *is orthogonal, i.e.* $SC_m \to SO(V)$ *for some invariant quadratic form on* V*. Then, when* $\mathcal{E}_\eta(V)$ *is a divisor it has multiplicity 2, i.e.* $\mathcal{E}_\eta(V) = 2\mathcal{D}_\eta(V)$ *for a divisor* $\mathcal{D}_\eta(V) \in |r\Xi_\eta + \Theta(V)|$.

Proof. We are here excluding the case $\mathcal{N}(m) \times P_\eta^-$ for odd m. Then we have already observed above that by corollary 4.7 and (21), $h^0(\widetilde{C}_\eta, L \otimes p^*E(\mathbf{C}^m))$ is *even* for all $(L, E) \in P_\eta^\pm \times \mathcal{M}^\pm(\mathrm{Spin}_m)$ in the remaining cases. The proposition then follows from the determinantal description of the functor Θ given in section 2. $\square$

Just as for the case $\eta = 0$, one now views the divisor $\mathcal{D}_\eta(V)$ as giving a rational map—or, more precisely, a *pair* of rational maps if m is even:

$$\begin{array}{rcl} f_\eta^\pm : \mathcal{N}(m) & \to & |m\Xi_\eta|_{P_\eta^\pm} \\ E & \mapsto & \{L \in P_\eta^\pm | H^0(\widetilde{C}_\eta, L \otimes p^*E(V)) \neq 0\}. \end{array}$$

By Riemann-Roch, Serre duality and the fact that $E(V)$ is self-dual, one sees again that $f_\eta^\pm(E)$ is a *symmetric* divisor. This means that each component of $\mathcal{N}(m)$ maps either into $|m\Xi_\eta|_{+,P_\eta^\pm} = \mathbf{P}H^0_+(P_\eta^\pm, m\Xi_\eta)$ or into $|m\Xi_\eta|_{-,P_\eta^\pm} = \mathbf{P}H^0_-(P_\eta^\pm, m\Xi_\eta)$; and the claim is:

8.6 Proposition. *When $V = \mathbf{C}^m$ is the standard orthogonal representation, for $m \geq 3$, each $f_\eta = f_\eta^+$ respects parity:*

$$f_\eta : \mathcal{M}^\pm(\mathrm{Spin}_m) \to |m\Xi_\eta|_{\pm,P_\eta};$$

and when m is even f_η^- respects parity:

$$f_\eta^- : \mathcal{M}^\pm(\mathrm{Spin}_m) \to |m\Xi_\eta|_{\pm,P_\eta^-}.$$

Proof. This uses proposition 4.6 in the same way as the proof of proposition 7.2. We suppose first that m is odd; choose $E \in \mathcal{N}(m)$ and a theta characteristic $L \in \vartheta(P_\eta)$. By (28) this means $L = p^*N = p^*(\eta \otimes N)$ for some theta characteristics N and $\eta \otimes N$ *of the same parity.* Then we note that

$$h^0(\widetilde{C}_\eta, L \otimes p^*E(\mathbf{C}^m)) = h^0(C, N \otimes E(\mathbf{C}^m)) + h^0(C, \eta \otimes N \otimes E(\mathbf{C}^m)),$$

and that (by 4.6) $h^0(C, N \otimes E(\mathbf{C}^m))$ is odd, and hence nonzero, provided $E \in \mathcal{M}(\mathrm{Spin}_m)$ and $L \in \vartheta^-(P_\eta)$ *or* $E \in \mathcal{M}^-(\mathrm{Spin}_m)$ and $L \in \vartheta^+(P_\eta)$. So by part 1 of lemma 7.3 it follows that f_η maps $\mathcal{M}^\pm(\mathrm{Spin}_m)$ into $|m\Xi_\eta|_{\pm,P_\eta}$ respectively.

If m is even then the same argument, using part 2 of lemma 7.3, shows that $\mathcal{M}^-(\mathrm{Spin}_m)$ maps under f_η into $|m\Xi_\eta|_{-,P_\eta}$; and maps under f_η^- into $|m\Xi_\eta|_{-,P_\eta^-}$ (see (29)).

Finally consider $\mathcal{M}(\mathrm{Spin}_m)$ for even m: for any theta characteristics $N, \eta \otimes N \in \vartheta(C)$, theorem 4.12 tells us that $h^0(C, N \otimes E(\mathbf{C}^m)) = h^0(C, \eta \otimes N \otimes E(\mathbf{C}^m)) = 0$ for generic $E \in \mathcal{M}(\mathrm{Spin}_m)$, so that generically $h^0(\widetilde{C}_\eta, L \otimes p^* E(\mathbf{C}^m)) = 0$ for any theta characteristic $L \in \vartheta(P_\eta)$ or $\vartheta(P^-_\eta)$. This implies (for each of P_η, P^-_η) that $\mathcal{M}(\mathrm{Spin}_m)$ does not map into $|m\Xi_\eta|_-$ and therefore maps into $|m\Xi_\eta|_+$ as required. □

As a consequence, pull-back of hyperplane sections under each $f^\pm_\eta$ is dual to a homomorphism (the analogue of (27))

$$s^\pm_\eta : H^0(\mathcal{N}(m), \Theta(V))^\vee \to H^0(P^\pm_\eta, r\Xi_\eta), \tag{31}$$

where $r = \dim V$, and where s^-_η is defined only for even m. Moreover, proposition 8.6 says that these homomorphisms *respect parity*—in other words the analogue of corollary 7.4 for this situation is:

8.7 Corollary. *When $V = \mathbf{C}^m$ is the standard orthogonal representation the homomorphisms $s^\pm_\eta$ respect parity:*

$$s_\eta = s^+_\eta : H^0(\mathcal{M}^\pm(\mathrm{Spin}_m), \Theta(\mathbf{C}^m))^\vee \to H^0_\pm(P_\eta, m\Xi_\eta) \qquad \textit{respectively},$$

and if m is even then additionally:

$$s^-_\eta : H^0(\mathcal{M}^\pm(\mathrm{Spin}_m), \Theta(\mathbf{C}^m))^\vee \to H^0_\pm(P^-_\eta, m\Xi_\eta) \qquad \textit{respectively}.$$

References

[ACGH] E. Arbarello, M. Cornalba, P.A. Griffiths, J. Harris: Geometry of Algebraic Curves, Springer 1985;

[A] M.F. Atiyah: Riemann surfaces and spin structures, Ann. Sci. Éc. Norm. Sup., 4 (1971) 47–62;

[B1] A. Beauville: Fibrés de rang 2 sur une courbe, fibré déterminant et fonctions theta, Bull. Soc. Math. France, 116 (1988) 431–448;

[B2] A. Beauville: Fibré de rang deux sur une courbe, fibré déterminant et fonctions theta II, Bull. Soc. Math. France, 119 (1991) 259–291;

[B3] A. Beauville: Conformal blocks, fusion rules and the Verlinde formula, preprint 1994;

[BL] A. Beauville, Y. Laszlo: Conformal blocks and generalised theta functions, Commun. Math. Phys. 164 (1994) 385–419;

[DN] J.-M. Drezet, M.S. Narasimhan: Groupe de Picard de variétés de modules de fibrés semi-stable sur les courbe algébriques, Invent. Math. 97 (1989) 53–94;

[F] G. Faltings: A proof for the Verlinde formula, J. Alg. Geometry 3 (1994) 347–374;

[vGP1] B. van Geemen, E. Previato: Prym varieties and the Verlinde formula, Math. Ann. 294 (1992) 741–754;

[vGP2] B. van Geemen, E. Previato: Heisenberg action and Verlinde formulas, in 'Integrable Systems, Verdier Memorial Conference' Luminy 1991, eds. O. Babelon et al, Birkhäuser (1994);

[LB] H. Lange, Ch. Birkenhake: Complex Abelian Varieties, Springer 1992;

[LS] Y. Laszlo, C. Sorger: The line bundles on the stack of parabolic G-bundles over curves and their sections, Institut de Mathématiques de Jussieu, preprint no. 40, 1995;

[KNR] S. Kumar, M.S. Narasimhan, A. Ramanathan: Infinite Grassmannian and moduli space of G-bundles, Math. Ann. 300 (1993) 395–423;

[M] D. Mumford: Theta-characteristics of an algebraic curve, Ann. Sci. Éc. Norm. Sup., 4 (1971) 181–192;

[N] M. Nagata: On self-intersection number of a section on a ruled surface, Nagoya Math. J. 37 (1970) 191–196;

[O1] W.M. Oxbury: Theta-characteristics and stable vector bundles, Proc. Lond. Math. Soc. (3) 71 (1995) 481–500;

[O2] W.M. Oxbury: Prym varieties and the moduli of spin bundles, to appear in proceedings of the Annual Europroj Conference, Barcelona 1994;

[OP] W.M. Oxbury, C. Pauly: $SU(2)$-Verlinde spaces as theta spaces on Pryms, to appear in Int. J. Math.;

[OW] W.M. Oxbury, S.M.J. Wilson: Reciprocity laws in the Verlinde formulae for the classical groups, to appear in Trans. Am. Math. Soc.;

[R1] S. Ramanan: Orthogonal and spin bundles over hyperelliptic curves, in 'Geometry and Analysis', papers dedicated to V.K. Patodi, Springer (1981);

[R2] S. Ramanan: The Schottky configuration, preprint 1995;

[Rth1] A. Ramanathan: Stable principal bundles on a compact Riemann surface, Math. Ann. 213 (1975) 129–152;

[Rth2] A. Ramanathan: Stable principal bundles on a compact Riemann surface: construction of moduli space, Bombay thesis, 1976;

[S] J.-P. Serre: Revêtements à ramification impaire et thêta-caractéristiques, C. R. Acad. Sci. Paris 311(I) (1990) 547–552;

[T] M. Thaddeus: Conformal field theory and the cohomology of the moduli space of stable bundles, J. Diff. Geometry 35 (1992) 131–149.

Department of Mathematical Sciences,
Science Laboratories,
South Road,
Durham DH1 3LE,
U.K.

E-mail: w.m.oxbury@durham.ac.uk